arXiv:1303.2274v1 [math.AG] 9 Mar 2013

REDUNDANT BLOW-UPS AND COX RINGS OF RATIONAL SURFACES

DONGSEON HWANG AND JINHYUNG PARK

Abstract. We introduce the notion of redundant blow-ups of rational surfaces based on Sakai's work, and show that having a finitely generated Cox ring is invariant under the redundant blow-up. As an application, we discuss the birational structure of rational surfaces with finitely generated Cox rings. In particular, we construct series of new examples of rational surfaces with anticanonical Iitaka dimension 0 and $-\infty$ of arbitrarily large Picard number whose Cox rings are finitely generated. This method is also applied to construct rational surfaces with big anticanonical divisor that are not minimal resolutions of del Pezzo surfaces containing at worst rational singularities.

Contents

1. Introduction 1
Acknowledgements 4
2. Theory of redundant blow-ups I: finite generation of Cox rings 4
3. Theory of redundant blow-ups II: redundant points 11
4. Rational surfaces with big anticanonical divisor 21
5. Minimal resolutions of rational $\mathbb{Q}$-homology projective planes 25
6. Examples of Mori dream rational surfaces with $\kappa(-K) \leq 0$ 28
References 29

1. Introduction

Throughout the paper, we work over an algebraically closed field k of arbitrary characteristic.

Algebraic varieties with finitely generated Cox rings, equivalently Mori dream spaces, have been received a lot of attention from various motivations. However, the classification problem of such varieties has been remained as a difficult problem even in the case of rational surfaces. Recently, it was shown that the Cox ring of every *big anticanonical rational surface*, i.e., a smooth projective rational surface with big anticanonical divisor, is finitely generated independently by Testa, Várilly-Alvarado and Velasco ([TVAV, Theorem 1]) and, Chen and Schnell ([CS, Theorem 3]). Motivated by this result, Artebani and Laface investigated a rational surface

---

Date: March 8, 2013.

2010 *Mathematics Subject Classification.* Primary 14J26; Secondary 14J17.

Key words and phrases. redundant blow-up, rational surface, Cox ring, Mori dream space, Zariski decomposition.

S with finitely generated Cox ring according to its *anticanonical Iitaka dimension*

$$\kappa(-K_S) := \max\{\dim \varphi_{|-nK_S|}(S) : n \in \mathbb{N}\}$$

whose value is one of $2, 1, 0$ and $-\infty$ ([AL]).

Let S be a smooth projective rational surface. If $\kappa(-K_S) = 2$, i.e., S is a big anticanonical rational surface, then the Cox ring of S is finitely generated as already mentioned. If $\kappa(-K_S) = 1$, then the Cox ring of S is finitely generated if and only if the relatively minimal model of the elliptic fibration $\pi\colon S \to \mathbb{P}^1$ has a Jacobian fibration with a finite Mordell-Weil group by [AL, Theorem 4.2, Theorem 4.8 and Corollary 5.4] in characteristic 0. On the contrary, very little is known about the classification when $\kappa(-K_S) \leq 0$. There have been some studies for a rational surface S with finitely generated Cox ring when $\kappa(-K_S) = 0$, and only one such example (that is a Coble rational surface) was recently given by Laface and Testa ([LT, Theorem 6.3]). However, as far as the authors know, no example has been known for the case $\kappa(-K_S) = -\infty$.

The principal aim of the present paper is to interpret this state of the art of the classification in terms of redundant blow-ups, motivated by Sakai's work on big anticanonical rational surfaces.

Let S be a smooth projective rational surface with $\kappa(-K_S) \geq 0$, and $-K_S = P + N$ be the Zariski decomposition. A point p in S is called a *redundant point* if $\operatorname{mult}_p N \geq 1$. The blow-up $f\colon \widetilde{S} \to S$ at a redundant point p is called a *redundant blow-up*. See Remark 2.4 for the original terminology by Sakai. The following is the first main theorem of this paper.

Theorem 1.1. *Let S be a smooth projective rational surface with $\kappa(-K_S) \geq 0$ and $f\colon \widetilde{S} \to S$ be a redundant blow-up at a point p in S. Assume that the Cox ring of S is finitely generated. Then the Cox ring of $\widetilde{S}$ is also finitely generated and $\kappa(-K_S) = \kappa(-K_{\widetilde{S}})$ unless $\kappa(-K_S) = 0$ and $\operatorname{mult}_p(N) = 1$ where $-K_S = P + N$ is the Zariski decomposition.*

Remark 1.2. The image of a surjective morphism from a Mori dream space is again a Mori dream space by [Ok, Theorem 1.1]. However, the converse does not hold in general. It is an interesting question to ask under what condition the converse is true.

Remark 1.3. There exist rational surfaces S with $\kappa(-K_S) = 0$ and $\operatorname{mult}_p(N) = 1$ whose Cox rings are finitely generated (see Remark 6.3). We do not know the finite generation of the Cox rings of $\widetilde{S}$ in this case.

The notion of redundant blow-up characterizes the morphisms between rational surfaces with finitely generated Cox rings of the same anticanonical Iitaka dimension $\kappa(-K) = 0$, which does not hold when $\kappa(-K) \neq 0$.

Theorem 1.4. *Let S be a smooth projective rational surface of $\kappa(-K_S) = 0$ with finitely generated Cox ring, and $f\colon \widetilde{S} \to S$ be a blow-up at a point. Suppose that the Cox ring of $\widetilde{S}$ is also finitely generated. Then $\kappa(-K_{\widetilde{S}}) = \kappa(-K_S)$ if and only if f is a redundant blow-up.*

Problem 1.5. *Classify all smooth projective rational surfaces of $\kappa(-K) = 0$ with finitely generated Cox rings from which every such surface can be obtained by a sequence of redundant blow-ups.*

Now, thanks to Theorem 1.1, we can construct many new rational surfaces with finitely generated Cox rings by taking redundant blow-ups. To achieve this goal, we only need to know the information of the redundant points on S.

Theorem 1.6. *Let $\bar{S}$ be a normal projective rational surface containing at worst rational singularities, and $\widetilde{S}$ be a surface obtained by a sequence of redundant blow-ups from the minimal resolution S of $\bar{S}$. Assume that $-K_{\bar{S}}$ is nef. Then there are only finitely many redundant points on $\widetilde{S}$ if and only if $\bar{S}$ contains at worst log terminal singularities. Moreover, the location of the redundant points is as follows:*

1. *If $\bar{S}$ contains at worst log terminal singularities, then every redundant point on $\widetilde{S}$ lies on the intersection points of the two curves contracted by the morphism $h\colon \widetilde{S} \to \bar{S}$.*
2. *If $\bar{S}$ contains a non-log terminal rational singularity, then there is a curve in $\widetilde{S}$ contracted by the morphism $h\colon \widetilde{S} \to \bar{S}$ such that every point lying on this curve is a redundant point.*

Moreover, the existence of the redundant points can be read off from the singularity types of $\bar{S}$.

Theorem 1.7. *Let $\bar{S}$ be a normal projective rational surface containing at worst rational singularities, and $g\colon S \to \bar{S}$ be its minimal resolution. Assume that $-K_{\bar{S}}$ is nef. Then we have the followings.*

1. *The number of rational surfaces obtained by a sequence of redundant blow-ups from S is finite if and only if $\bar{S}$ contains at worst log terminal singularities.*
2. *S has no redundant point if and only if $\bar{S}$ contains at worst canonical singularities or log terminal singularities whose dual graphs are as follows:*



Now we are ready to construct series of infinitely many new examples of rational surfaces with finitely generated Cox rings for $\kappa(-K_S) = 0$ and $-\infty$ using redundant blow-ups.

Theorem 1.8. *For each $n \geq 11$, there exist smooth projective rational surfaces S and $\widetilde{S}$ with finitely generated Cox rings such that*

1. *$\kappa(-K_S) = 0$ and $\rho(S) = n$.*
2. *$\kappa(-K_{\widetilde{S}}) = -\infty$ and $\rho(\widetilde{S}) = n + 1$.*

where ρ denotes the Picard number.

The theorem is proven by explicit construction (see Example 6.1). In fact, they are obtained by blow-ups from extremal rational elliptic surfaces with effective k^*-actions. By a result of Knop ([Kn]), all the rational surfaces with effective k^*-actions have finitely generated Cox rings. There are four extremal rational elliptic surfaces having effective $\mathbb{C}^*$-action (see Remark 6.2), and their Cox rings are explicitly calculated in [AGL] based on the technique developed in [HS, Theorem 1.3].

The redundant blow-up method is also applied to construct big anticanonical rational surfaces that are not minimal resolutions of del Pezzo surfaces containing at worst rational singularities, which provides a negative answer to a question remarked in [TVAV, Remark 3].

Theorem 1.9. *For each $n \geq 10$, there exists a big anticanonical rational surface with Picard number n that is not a minimal resolution of a del Pezzo surface containing at worst rational singularities.*

Finally, we propose another view point on rational surfaces with finitely generated Cox rings. Recall that the Cox ring of the surface obtained by blowing up at least 9 very general points on the projective plane is not finitely generated. However, the Cox ring of the surface obtained by blowing up at any number of points on a line on the projective plane is always finitely generated ([EKW, Example 3.3] and [Ot]). Thus we observe that a rational surface obtained by blowing up at points in special position has a tendency to have a finitely generated Cox ring. This observation motivates us to study the extremal case: finite generation of the Cox ring of the minimal resolution of a rational surface with Picard number one.

Theorem 1.10. *Let $\bar{S}$ be a normal projective rational surface of Picard number one with at worst log terminal singularities, and $g\colon S \to \bar{S}$ be its minimal resolution. Assume that $-K_{\bar{S}}$ is nef and $\bar{S}$ does not contain canonical singularities. Then the Cox ring of S is finitely generated.*

In the case of $k = \mathbb{C}$, a normal projective surface with at worst quotient singularities and the second topological Betti number $b_2 = 1$ is called a $\mathbb{Q}$-*homology projective plane.* In [HwK3], one can find examples of $\mathbb{Q}$-homology projective planes satisfying the condition of Theorem 1.10. From them, we can obtain more examples of rational surfaces of $\kappa(-K) = 0$ with finitely generated Cox rings that are not Coble rational surfaces (see Remark 5.5). On the other hand, there exist infinitely many $\mathbb{Q}$-homology projective planes which are rational surfaces with ample canonical divisors (see [HwK3]). However, we do not know the finite generation of their Cox rings.

Question 1.11. *Let $\bar{S}$ be a $\mathbb{Q}$-homology projective plane, and S be its minimal resolution. Is the Cox ring of S finitely generated?*

The paper is organized as follows. We develop the theory of redundant blow-ups and prove Theorem 1.1 and Theorem 1.4 in Section 2. Section 3 is devoted to investigate redundant points. As applications, we provide examples of big anticanonical rational surfaces that are not minimal resolutions of del Pezzo surfaces with rational singularities and study weak del Pezzo surfaces in Section 4, consider the finite generation of Cox rings of minimal resolutions of $\mathbb{Q}$-homology projective planes in Section 5 and finally construct infinitely many new rational surfaces with finitely genrated Cox rings with anticanonical Iitaka dimension 0 as well as $-\infty$ in Section 6.

Acknowledgements

The authors would like to thank Junmyeong Jang for useful comments on the techniques in positive characteristic.

2. Theory of redundant blow-ups I: finite generation of Cox rings

In this section, we develop the basic theory of redundant blow-ups, and prove Theorem 1.1 and Theorem 1.4.

2.1. **Basic setup.** Let S be a smooth projective surface, and D be a $\mathbb{Q}$-divisor. The *Iitaka dimension* of D is given by

$$\kappa(D) := \max\{\dim \varphi_{|-nD|(S)} : n \in \mathbb{N}\}$$

whose value is one of $2, 1, 0$ and $-\infty$. We call $\kappa(-K_S)$ the *anticanonical Iitaka dimension* of S. Note that $\kappa(-K_S) \geq 0$ if and only if $-K_S$ is pseudo-effective ([Sa, Lemma 3.1]).

Let D be a pseudo-effective divisor. We will frequently use the notion of the *Zariski decomposition* of D (see [Sa, Section 2] for details): D can be written uniquely as $P + N$, where P is a nef $\mathbb{Q}$-divisor, N is an effective $\mathbb{Q}$-divisor, $P.N = 0$, and the intersection matrix of irreducible components of N is negative definite.

Let $D = \sum_{i=1}^{n} \alpha_i E_i$ be an effective $\mathbb{Q}$-divisor on S where E_i is a prime divisor for all $1 \leq i \leq n$. The *multiplicity* of D at a point p in S is defined by $\operatorname{mult}_p(D) := \sum_{i=1}^{n} \alpha_i \operatorname{mult}_p(E_i)$ where $\operatorname{mult}_p(E_i)$ denotes the usual multiplicity of E_i at p.

Definition 2.1. Let S be a smooth projective rational surface with $\kappa(-K_S) \geq 0$, and $-K_S = P + N$ be the Zariski decomposition. A point p in S is called a *redundant point* if $\operatorname{mult}_p N \geq 1$. The blow-up $\widetilde{S} \to S$ at a redundant point p is called a *redundant blow-up*, and the exceptional curve E is called a *redundant curve*.

Remark 2.2. By [Sa, Lemma 2.4], N is in the fixed part of $|-K_S|$, so $\operatorname{mult}_p N$ is well-defined.

Now we reformulate Sakai's result on redundant blow-ups. It will play a key role in the whole paper.

Lemma 2.3 (Corollary 6.7 and Lemma 6.9 of [Sa]). *Let S be a smooth projective rational surface with $\kappa(-K_S) \geq 0$, and $f\colon \widetilde{S} \to S$ be a blow-up at a point p with the exceptional divisor E. Assume that $\kappa(-K_{\widetilde{S}}) \geq 0$. Then f is a redundant blow-up if and only if $\widetilde{P} = f^*P$ and $\widetilde{N} = f^*N - E$ where $-K_S = P + N$ and $-K_{\widetilde{S}} = \widetilde{P} + \widetilde{N}$ are the Zariski decompositions. Moreover, if f is a redundant blow-up, then $\kappa(-K_{\widetilde{S}}) = \kappa(-K_S)$, and the intersection matrix of irreducible components of $f^*(N)$ is negative definite.*

Note that for a blow-up $f\colon \widetilde{S} \to S$, we have $\kappa(-K_S) \geq \kappa(-K_{\widetilde{S}})$ in general.

Remark 2.4. In Lemma 2.3, since $\widetilde{P}.E = 0$, the exceptional curve E is redundant in the sense of Sakai ([Sa, Definition 4.1]). Sakai defined the notion of redundant curve only for big anticanonical rational surfaces. Note that, when both $\widetilde{X}$ and X are big anticanonical rational surfaces, the two notions coincide.

Theorem 2.5 (Proposition 2.9 of [HuK] and Corollary 1 of [GM]). *Let S be a smooth projective surface with $q(S) = 0$. Then the Cox ring of S is finitely generated if and only if*

1. *the effective cone* $\operatorname{Eff}(S)$ *is rational polyhedral, and*
2. *every nef divisor is semiample.*

Such a surface is called a *Mori dream surface*. The following lemma will be useful in proving that the effective cone is rational polyhedral. We remark that the proof in [AL] works for arbitrary characteristic, since the cone theorem still holds for positive characteristic by [Ta, Theorem 4.4].

Lemma 2.6 (Corollary 2.2 of [AL]). *Let S be a smooth projective rational surface with $\kappa(-K_S) \geq 0$. Then $\mathrm{Eff}(S)$ is rational polyhedral if and only if there are only finitely many (-1)-curves and (-2)-curves on S.*

2.2. **Finite generation of Cox rings.** In this subsection, we give a proof of Theorem 1.1. First, we prove some lemmas (cf. [AL, Proposition 1.3] and [AL, Lemma 4.7]).

Lemma 2.7. *Let S be a smooth projective surface with $q(S) = 0$, and M be an effective base point free $\mathbb{Z}$-divisor on S such that $M^2 = 0$ and $M.K_S = 0$. If M is nontrivial, then the linear system $|M|$ induces an elliptic (or quasi-elliptic) fibration $\varphi_{|M|}\colon S \to \mathbb{P}^1$.*

Proof. We slightly modify [AL, Proof of Proposition 1.3] to make it work for arbitrary characteristic. Since M is not big and is base point free, we have $\kappa(M) = 1$. Thus, the morphism $\varphi_{|M|}\colon S \to B \subset \mathbb{P}^n$ maps to an integral curve. Consider the Stein factorization

$$\begin{tikzcd} S \arrow[rr, "\varphi_{|M|}"] \arrow[dr, "f"'] & & B \\ & B' \arrow[ur, "g"'] & \end{tikzcd}$$

where B' is an integral curve, f is a contraction, and g is a finite morphism. We have the Leray spectral sequence

$$E_2^{p,q} = H^p(B', R^q f_* \mathcal{O}_S) \Rightarrow H^{p+q}(S, \mathcal{O}_S).$$

Since the edge homomorphism $H^1(B', \mathcal{O}_{B'}) = E_2^{1,0} \to H^1(S, \mathcal{O}_S)$ is injective, we obtain $0 = q(S) \geq p_a(B')$. Thus, $B' \simeq \mathbb{P}^1$. Let $a := \deg(g) \deg_{\mathbb{P}^n}(B)$. Then $M = aD$ where $\mathcal{O}_S(D) = f^*\mathcal{O}_{\mathbb{P}^1}(1)$. Since $h^0(D) \geq 2$, we have

$$n + 1 = h^0(M) = h^0(aD) \geq a + 1.$$

On the other hand, since the embedding $B \subset \mathbb{P}^n$ is non-degenerate, $a \geq \deg_{\mathbb{P}^n}(B) \geq n$. Thus, $a = \deg_{\mathbb{P}^n}(B) = n$, so B is a curve of minimal degree. By the classification of varieties of minimal degree (see [EH, Theorem 1]), we conclude that g is a Veronese embedding, and hence we get a fibration $\varphi_{|M|}\colon S \to B \simeq \mathbb{P}^1$ with connected fibers. Note that the condition $(\varphi_{|M|})_*\mathcal{O}_S = \mathcal{O}_{\mathbb{P}^1}$ is equivalent to the function field $k(\mathbb{P}^1)$ being algebraically closed in $k(S)$. By [Ba, Corollary 7.3], all but finitely many fibers of $\varphi_{|M|}$ are integral curves. Now, the assertion immediately follows from the adjunction formula. □

The following lemma is the technical main ingredient of this subsection.

Lemma 2.8. *Let S be a Mori dream surface. Then, for any fixed integer $r \geq 0$, there is a finite number of classes $[C] \in \mathrm{Eff}(S)$ such that $C^2 = r^2 - 1, K_S.C = -r-1$ and C is an integral curve.*

Proof. Since S is a Mori dream surface, there are only finitely many (-1)-curves. Thus, we assume that $r \geq 1$. It suffices to show that every class $[C]$ lies in some bounded region in the Néron-Severi space $N^1(S)_{\mathbb{R}} := N^1(S) \otimes \mathbb{R}$, because $[C]$ is an integral point.

If $r = 1$, then $C^2 = 0$ and $K_S.C = -2$, so C is a nef divisor on S. It is easy to check the assertion when the Picard number $\rho(S) \leq 2$. Thus, assume that $\rho(S) \geq 3$. Then, by [AL, Proposition 1.1] or [Ni, p.75], $\mathrm{Eff}(S)$ is generated by the classes of integral curves with negative self-intersection. Since $[C] \in \mathrm{Eff}(S)_{.K_S<0}$, we may write

$$C = \sum_{i=1}^{n} e_i E_i + \sum_{j=1}^{k} f_j F_j$$

where every $e_i, f_j \in \mathbb{Q}_{>0}$, each E_i is a (-1)-curve, and each F_j is a (-2)-curve. Since there are only finitely many (-1)-curves and (-2)-curves on S, it suffices to show that all e_i and f_j are bounded.

We have

$$2 = C.(-K_S) = \sum e_i E_i.(-K_S) + \sum f_j F_j.(-K_S) = \sum e_i,$$

so each $e_i \leq 2$, i.e., every e_i is bounded. Now, we have

$$0 = C^2 = \sum e_i(E_i.C) + \sum f_j(F_j.C),$$

so $E_i.C = 0$ and $F_j.C = 0$ for every i and j.

To derived a contradiction, suppose that $f_j > 2^{k+1}$ for some j. By reordering the indices, there exists a positive integer $u < k$ such that for some positive integer $m \leq k$, we have $f_{j_1} \leq 2^m$ for every j_1 with $1 \leq j_1 \leq u$, and $f_{j_2} > 2^{m+1}$ for every j_2 with $u+1 \leq j_2 \leq k$.

Claim 2.9. $F_{j_1}.F_{j_2} = 0$ for every j_1 and j_2 with $1 \leq j_1 \leq u$ and $u+1 \leq j_2 \leq k$.

Proof. If $F_{j_1}.F_{j_2} \geq 1$ for some j_1 and j_2 with $1 \leq j_1 \leq u$ and $u+1 \leq j_2 \leq k$, then we have

$$0 = F_{j_1}.C = \sum e_i F_{j_1}.E_i + \sum f_j F_{j_1}.F_j,$$

so we obtain

$$2^{m+1} \geq 2f_{j_1} = \sum_i e_i F_{j_1}.E_i + \sum_{j \neq j_1} f_j F_{j_1}.F_j \geq f_{j_2} > 2^{m+1},$$

which is a contradiction. □

Claim 2.10. $E_i.F_{j_2} = 0$ for every i and j_2 with $u+1 \leq j_2 \leq k$.

Proof. If $E_{i_0}.F_{j_2} \geq 1$ for some i_0 and j_2 with $u+1 \leq j_2 \leq k$, then we have

$$0 = E_{i_0}.C = \sum e_i(E_{i_0}.E_i) + \sum f_j(E_{i_0}.F_j),$$

so we obtain

$$2 \geq e_{i_0} = \sum_{i \neq i_0} e_i(E_{i_0}.E_i) + \sum_j f_j(E_{i_0}.F_j) \geq f_{j_2} > 2^{m+1},$$

which is a contradiction. □

Denote $D' := \sum_{i=1}^{n} e_i E_i + \sum_{j_1=1}^{u} f_{j_1} F_{j_1}$ and $D := C - D' = \sum_{j_2=u+1}^{l} f_{j_2} F_{j_2}$. Note that $C.D = \sum_{j_2=u+1}^{l} f_{j_2} C.F_{j_2} = 0$. By Claim 2.9 and 2.10, we have $D.E_i = 0$ and $D.F_{j_1} = 0$ for every i and j_1 with $1 \leq j_1 \leq u$. Thus, we have $D'.D = 0$, so we obtain $D^2 = 0$ by the following calculation:

$$0 = C.D = (D + D').D = D^2 + D'.D = D^2.$$

On the other hand, we claim that D is nef. Indeed, for every j_2 with $u+1 \leq j_2 \leq k$, by Claim 2.9 and 2.10, we have $D'.F_{j_2} = 0$, so we obtain $D.F_{j_2} = 0$ by the following calculation:

$$0 = C.F_{j_2} = D.F_{j_2} + D'.F_{j_2} = D.F_{j_2}.$$

Since S is a Mori dream surface, D is semiample. Note that F_{j_2} is (-2)-curve for every j_2 with $u+1 \leq j_2 \leq l$, so $D.K_S = 0$. By Lemma 2.7, for some integer $a > 0$, the linear system $|aD|$ gives a fibration $S \to \mathbb{P}^1$. Since $C.D = 0$, the integral curve C is contained in a fiber S_s of the fibration $S \to \mathbb{P}^1$. Since $C^2 = 0$, by the Zariski lemma on fibration (see [Ba, Corollary 2.6]), $C = rS_s$ for some $r \in \mathbb{Q}$. It is a contradiction, because $S_s.K_S = 0$ but $C.K_S = -2$. Hence, every f_j is bounded. We complete the proof for the case $r = 1$.

Finally, we consider the remaining case $r \geq 2$. In this case, $C^2 = r^2 - 1 > 0$, so C is nef and big. Let $A_1, \ldots, A_k$ be nontrivial $\mathbb{Z}$-divisors generating the nef cone $\operatorname{Nef}(S)$. We may write $C = \sum_i a_i A_i$ where $a_i \in \mathbb{Q}_{\geq 0}$ for every i. By the Hodge index theorem, $A_i.C \geq 1$ for every i. We have

$$r^2 - 1 = C^2 = \sum a_i A_i.C \geq \sum a_i,$$

so every a_i is bounded. □

Proof of Theorem 1.1. Let S be a Mori dream rational surface and $f\colon \widetilde{S} \to S$ be a redundant blow-up at a point p with the exceptional divisor E. Then $\widetilde{S}$ is a rational surface with $\kappa(-K_S) = \kappa(-K_{\widetilde{S}})$ by Lemma 2.3. Let $-K_{\widetilde{S}} = \widetilde{P} + \widetilde{N}$ and $-K_S = P + N$ be the Zariski decompositions. By Lemma 2.3, we have $\widetilde{P} = f^*P$ and $\widetilde{N} = f^*N - E$. Note that every integral curve on $\widetilde{S}$ different from E can be written as $\widetilde{C} = f^*C - rE$ where C is an integral curve on S and $r \geq 0$ is an integer. Note that $C^2 = r^2 - 1$ and $K_S.C = -r - 1$. Now we have three cases.

Case 1: $\kappa(-K_S) = 2$.

By [TVAV, Theorem 1], $\widetilde{S}$ is a Mori dream rational surface.

Case 2: $\kappa(-K_S) = 1$.

By Lemma 2.7, for some integer $m > 0$, the linear system $|m\widetilde{P}|$ induces a fibration $\varphi\colon \widetilde{S} \to \mathbb{P}^1$. First, we show that the effective cone $\operatorname{Eff}(\widetilde{S})$ is rational polyhedral. By Lemma 2.6, it is sufficient to show that there are only finitely many (-1)-curves and (-2)-curves on $\widetilde{S}$. Let $\widetilde{C}$ be a (-2)-curve. Since $0 = -K_{\widetilde{S}}.\widetilde{C} = \widetilde{P}.C + \widetilde{N}.C$, the curve $\widetilde{C}$ is contained in either a reducible fiber of φ or the support of $\widetilde{N}$. Thus, $\widetilde{S}$ contains finitely many (-2)-curves.

Let $\widetilde{C}$ be a (-1)-curve on $\widetilde{S}$ different from E. In this case, we use an idea in [AL, Proof of Theorem 4.8]. By [Sa, Theorem 3.4], there exists a birational morphism $g\colon S \to S_0$ such that S_0 is a relatively minimal model of the fibration $\varphi_{|P|}\colon S \to \mathbb{P}^1$. Then $-K_{S_0} = P_0$ is the Zariski decomposition. Since S has a redundant point, $S \neq S_0$. Moreover, by [Sa, Theorem 3.4] or [AL$'$, Lemma 4.4] (see also Lemma 2.13), $P = \alpha g^* P_0$ where α is a rational number with $0 < \alpha < 1$. Since $-K_{\widetilde{S}}.\widetilde{C} = 1$, either $\widetilde{C}$ is a component of N or a reducible fiber of φ, or $0 < \widetilde{C}.\widetilde{P} \leq 1$. For the first two cases, there are only finitely many possibilities for such a (-1)-curve $\widetilde{C}$. Assume the third case. We further assume that $\widetilde{C}$ is not a component of the exceptional divisor of $g \circ f\colon \widetilde{S} \to S_0$. Since $\widetilde{P} = \alpha f^* g^* P_0$, we have

$$r = \widetilde{C}.E \leq \widetilde{C}.(f^*g^*(-K_{S_0}) + K_{\widetilde{S}}) = \frac{1}{\alpha}\widetilde{C}.\widetilde{P} - 1 \leq \frac{1}{\alpha} - 1,$$

i.e., r is bounded. Now, Lemma 2.8 implies that $\widetilde{S}$ contains only finitely many (-1)-curves. Indeed, if $f^*C - rE$ and $f^*C' - rE$ are (-1)-curves on $\widetilde{S}$ for two distinct linearly equivalent integral curves C and C' on S, then $f^*C - rE$ and

$f^*C' - rE$ are also linearly equivalent. So $f^*C - rE = f^*C' - rE$. Thus, at most one integral curve in each class $[C]$ gives a (-1)-curve in $\widetilde{S}$.

Since the proof of [AL, Theorem 3.4] works for arbitrary characteristic, every nef divisor on $\widetilde{S}$ is semiample, and hence $\widetilde{S}$ is a Mori dream rational surface.

Case 3: $\kappa(-K_S) = 0$.

For this case, we have $P = 0$ since P is semiample and $\kappa(P) = 0$. Recall that $\widetilde{P} = f^*P = 0$ and hence $-K_{\widetilde{S}} = \widetilde{N} = f^*N - E$. By the assumption $m := \operatorname{mult}_p(N) > 1$, we have

$$\widetilde{N} = f_*^{-1}N + (m-1)E,$$

i.e., the redundant curve E is contained in the support of $\widetilde{N}$.

First, we show that the effective cone $\operatorname{Eff}(\widetilde{S})$ is rational polyhedral. By Lemma 2.6, it is sufficient to show that there are only finitely many (-1)-curves and (-2)-curves on $\widetilde{S}$. Let $\widetilde{C} = f^*C - rE$ be a (-2)-curve not in the support of $\widetilde{N}$. Then we have

$$0 = -K_{\widetilde{S}}.\widetilde{C} = \widetilde{N}.\widetilde{C} = (f_*^{-1}N + (m-1)E).\widetilde{C} \geq (m-1)E.\widetilde{C},$$

so $E.\widetilde{C} = 0$. Since $E.\widetilde{C} = r$, we obtain $r = 0$, i.e., C is a (-2)-curve. Thus, there are finitely many (-2)-curves on $\widetilde{S}$.

Let $\widetilde{C} = f^*C - rE$ be a (-1)-curve not in the support of $\widetilde{N}$. Then we have

$$1 = -K_{\widetilde{S}}.\widetilde{C} = \widetilde{N}.\widetilde{C} = (f_*^{-1}N + (m-1)E).\widetilde{C} \geq (m-1)E.\widetilde{C} = (m-1)r,$$

so $r \leq \frac{1}{m-1}$, i.e., r is bounded. By Lemma 2.8, there are only finitely many (-1)-curves on $\widetilde{S}$.

Finally, we show that every nef divisor $\widetilde{M}$ on $\widetilde{S}$ is semiample. By [LT, Lemma 3.1], any nef divisor $\widetilde{M}$ with $-K_{\widetilde{S}}.\widetilde{M} > 0$ is semiample. Since $-K_{\widetilde{S}}$ is effective, we may assume that $-K_{\widetilde{S}}.\widetilde{M} = 0$. We have

$$0 = -K_{\widetilde{S}}.\widetilde{M} = \widetilde{N}.\widetilde{M} = (f_*^{-1}N + (m-1)E).\widetilde{M} \geq (m-1)E.\widetilde{M},$$

so we obtain $E.\widetilde{M} = 0$. We may write $\widetilde{M} = f^*M - rE$ for some divisor M on S. Then $r = E.\widetilde{M} = 0$. For any effective curve C on S, we have

$$M.C = f^*M.f^*C = \widetilde{M}.f^*C \geq 0,$$

so M is also nef on a Mori dream surface S, hence it is semiample. Thus, $\widetilde{M} = f^*M$ is also semiample. ☐

Remark 2.11. It plays a crucial role in the proof that the exceptional curve E appears in the support of $\widetilde{N}$ when $\kappa(-K_S) = 0$.

2.3. **Classification of Mori dream rational surfaces.** Now, we investigate the classification problem of Mori dream rational surfaces with nonnegative anticanonical Iitaka dimension using redundant blow-ups, and we show Theorem 1.4. Let S be a smooth projective rational surface with $\kappa(-K_S) \geq 0$.

2.3.1. $\kappa(-K_S) = 2$. Recall that S is always a Mori dream rational surface by [TVAV, Theorem 1] or [CS, Theorem 3]. Moreover, we have the following classification result.

Theorem 2.12 (Proposition 4.2 and Theorem 4.3 of [Sa])**.** *Every big anticanonical rational surface can be obtained by a sequence of redundant blow-ups from the minimal resolution of a del Pezzo surface with rational singularities.*

We will further discuss the properties of big anticanonical rational surfaces in Section 4.

2.3.2. $\kappa(-K_S) = 1$. In the case of characteristic zero, we have the classification result as follows. By [AL′, Theorem 3.4 and Theorem 4.8], every Mori dream rational surface with anticanonical Iitaka dimension 1 can be obtained by a sequence of blow-ups of type (1) or (2) in Lemma 2.13 from a relatively minimal rational elliptic surface whose Jacobian fibration has a finite Mordell-Weil group. Note that using the Ogg-Shafarevich theory, every such rational elliptic surface can be obtained from extremal rational elliptic surface, which are completely classified in [MP]. We remark that quasi-elliptic fibration exists in characteristic 2 and 3.

We note that the proof in [AL′] works for arbitrary characteristic.

Lemma 2.13 (Lemma 4.4 of [AL′])**.** *Let $f\colon \widetilde{S} \to S$ be the blow-up at a point p in a smooth projective rational surface S with $\kappa(-K_S) = 1$. Then $\kappa(-K_{\widetilde{S}}) = \kappa(-K_S)$ if and only if one of the followings hold:*

(1) *f is a redundant blow-up, or*

(2) $\mathrm{mult}_p(N) < 1$ *but* $\mathrm{mult}_p(-K_S) > 1$.

For the second case, we have $\widetilde{P} = \frac{\mathrm{mult}_p(-K_X)-1}{\mathrm{mult}_p(P)} f^*P$ and $\widetilde{N} = \frac{1-\mathrm{mult}_p(N)}{\mathrm{mult}_p(P)} f^*P + f^*N - E$ where $-K_{\widetilde{S}} = \widetilde{P} + \widetilde{N}$ and $-K_S = P + N$ are the Zariski decompositions, and E is the exceptional divisor of f.

See [AL′, Definition 4.3] for well-definedness of $\mathrm{mult}_p(P)$ and $\mathrm{mult}_p(-K_S)$. The Mori dream rational surface S with $\kappa(-K_S) = 1$ in Example 6.1 admits infinitely many redundant blow-ups. In [AL′, Proposition 4.6], Artebani and Laface constructed an infinite sequence of blow-ups of type (2) in Lemma 2.13.

Furthermore, we can also prove that the blow-ups of type (1) and (2) of Lemma 2.13 preserves the finite generation of Cox ring (cf. [AL, Theorem 4.8]).

Proposition 2.14. *Let S be a Mori dream rational surface with $\kappa(-K_S) = 1$, and $f\colon \widetilde{S} \to S$ be the blow-up at a point p. If $\kappa(-K_{\widetilde{S}}) = \kappa(-K_S)$, then $\widetilde{S}$ is also a Mori dream rational surface.*

Proof. By Lemma 2.13, either f is a redundant blow-up, or $\mathrm{mult}_p(N) < 1$ but $\mathrm{mult}_p(-K_S) \geq 1$. For the first case, the assertion follows from Theorem 1.1. The proof of Theorem 1.1 for $\kappa(-K_S) = 1$ also can be applied to the remaining case. □

2.3.3. $\kappa(-K_S) = 0$. We do not have any classification result of Mori dream rational surface with anticanonical Iitaka dimension 0 at all, but we have a characterization of blow-ups preserving the finite generation of Cox rings.

Proof of Theorem 1.4. If f is a redundant blow-up, then $\kappa(-K_{\widetilde{X}}) = 0$ by Lemma 2.3. Conversely, assume that $\kappa(-K_{\widetilde{S}}) = 0$. Let $-K_S = P + N$ be the Zariski decomposition. Since $\kappa(-K_S) = \kappa(P) = 0$ and the nef divisor P on a Mori dream rational surface S is semiample, we have $P = 0$. Let E be the exceptional divisor of f. Then, for the Zariski decomposition $-K_{\widetilde{S}} = \widetilde{P} + \widetilde{N}$, we have $\widetilde{P} = 0$ and $\widetilde{N} = -K_{\widetilde{S}} = f^*(N) - E$. By Lemma 2.3, f is a redundant blow-up. □

Toward the classification, we construct many new examples of such surfaces and redundant blow-ups (see Section 5 and 6).

We close this section by asking the following.

Question 2.15. *Do analogous statements of Theorem 1.1 and 1.4 exist without the assumption* $\kappa(-K_X) \geq 0$*?*

3. Theory of redundant blow-ups II: redundant points

In this section, we focus on redundant points on rational surfaces, and prove Theorem 1.6 and Theorem 1.7. The main ingredients of the proofs are the Zariski decomposition and explicit calculation of discrepancies. In what follows, $\bar{S}$ denotes a normal projective rational surface containing at worst rational singularities, and $g\colon S \to \bar{S}$ denotes the minimal resolution. Throughout the section, we assume that $-K_{\bar{S}}$ is nef.

3.1. **Redundant points with respect to a given singularity.**

In this subsection, we investigate redundant points on S with respect to singularities of $\bar{S}$. We give a proof of Theorem 1.6 at the end.

First, we recall some basic facts concerning rational singularities on surfaces. A basic reference is Chapter 4 of [KM]. If an algebraic surface contains at worst rational singularities, then its singularities are isolated and the surface is $\mathbb{Q}$-factorial (see [Ba, Theorem 4.6]).

Let $(\bar{S}, s)$ be a germ of a rational singularity on a normal surface, and $g\colon S \to \bar{S}$ be the minimal resolution. Denote the exceptional set by $A = \pi^{-1}(s) = E_1 \cup \cdots \cup E_l$ where each E_i is an irreducible component. Note that $E_i^2 = -n_i \leq -2$ where each n_i is an integer for $i = 1, \ldots, l$ and A has a simple normal crossing (*snc* for short) support. Now we have

$$-K_S = g^*(-K_{\bar{S}}) + \sum_{i=1}^{l} a_i E_i$$

where each a_i is a nonnegative rational number for all $i = 1, \ldots, l$. Here, we call a_i the *discrepancy* of E_i with respect to $\bar{S}$ for the convenience in computation, even though $-a_i$ is called the discrepancy in the literature. This number can be obtained by the simultaneous linear equations:

$$\sum_{j=1}^{l} a_j E_j E_i = -K_S . E_i = -n_i + 2 \text{ for } i = 1, \ldots, l.$$

The second equality follows from the adjunction formula. Thus, each discrepancy a_i can be calculated by the following matrix equation

$$\begin{pmatrix} -n_1 & E_2E_1 & \cdots & E_lE_1 \\ E_2E_1 & -n_2 & \cdots & E_lE_2 \\ \vdots & \vdots & \ddots & \vdots \\ E_lE_1 & E_lE_2 & \cdots & -n_l \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_l \end{pmatrix} = - \begin{pmatrix} n_1 - 2 \\ n_2 - 2 \\ \vdots \\ n_l - 2 \end{pmatrix}.$$

We will often use the below well-known characterization of surface singularities, which can be regarded as a definition.

Remark 3.1. (1) A singularity s is *canonical* if and only if $a_i = 0$ for all $i = 1, \ldots, l$.
(2) A singularity s is *log terminal* if and only if $0 \leq a_i < 1$ for all $i = 1, \ldots, l$.

We note that every log terminal singularity is rational (see [KM, Theorem 4.12]). When $\mathrm{char}(k) = 0$, a log terminal singularity is nothing but a quotient singularity (see [KM, Proposition 4.18]).

Recall that $-K_{\bar{S}}$ is nef. Then we can get the Zariski decomposition of $-K_S$ immediately.

Lemma 3.2. *Let $P = g^*(-K_{\bar{S}})$ and $N = \sum_{i=1}^{l} a_i E_i$ where each E_i is a g-exceptional curve and each a_i is the discrepancy of E_i with respect to $\bar{S}$ for $1 \le i \le l$. Then $-K_S = P + N$ is the Zariski decomposition.*

Proof. The pull-back of a nef divisor is again nef. Clearly $P.N = 0$ and the intersection matrix of irreducible components of N is negative definite. □

Suppose that S contains a redundant point p. Let $f\colon \widetilde{S} \to S$ be the redundant blow-up at p and E be the exceptional divisor of f. Let $-K_{\widetilde{S}} = \widetilde{P} + \widetilde{N}$ be the Zariski decomposition. Then we obtain

$$\widetilde{N} = \sum_{i=1}^{l} a_i \widetilde{E}_i + (\mathrm{mult}_p(N) - 1)E = \sum_{i=1}^{l} a_i \widetilde{E}_i + \left(\sum_{p \in E_j} a_j - 1 \right) E$$

where $\widetilde{E}_i$ is the strict transform of E_i for each $1 \le i \le l$. We note that $\sum_{p \in E_j} a_j \ge 1$, since $\mathrm{mult}_p(N) \ge 1$.

Remark 3.3. Clearly N has a snc support, so is $\widetilde{N}$. Thus, the negative part of the Zariski decomposition of the anticanonical divisor of a big anticanonical rational surface has also a snc support by Theorem 2.12. This is not true anymore for Mori dream rational surfaces with anticanonical Iitaka dimension 1 or 0. E.g., some blow-up of the complex extremal rational elliptic surface X_{22} in [MP, Theorem 4.1] (see Remark 6.2).

In the rest of this subsection, we will see how to completely determine the location of redundant points on S in terms of the singularities of $\bar{S}$.

3.1.1. *$\bar{S}$ contains at worst canonical singularities.*

Lemma 3.4. *If $\bar{S}$ has at worst canonical singularities, then S has no redundant points.*

Proof. By Remark 3.1, $N = 0$, thus the lemma follows. □

3.1.2. *$\bar{S}$ contains at worst log terminal singularities.*

Lemma 3.5. *If $\bar{S}$ contains at worst log terminal singularities, then any surface $\widetilde{S}'$ obtained by a sequence of redundant blow-ups from S has finitely many redundant points, and every redundant point is an intersection point of two curves contracted by the morphism $\widetilde{S}' \to \bar{S}$.*

Proof. By Remark 3.1, $0 \le a_i < 1$ for each $1 \le i \le l$. Finding a point p in S satisfying $\mathrm{mult}_p(N) \ge 1$ is equivalent to finding an intersection point of E_j and E_k such that $a_j + a_k \ge 1$ for some $1 \le j \le l$ and $1 \le k \le l$. Since the number of E_i's is finite, the number of redundant points is also finite. In this case, we have

$$\widetilde{N} = \sum_{i=1}^{l} a_i \widetilde{E}_i + (a_j + a_k - 1)E.$$

Observe that $a_i < 1$ for each $1 \leq i \leq l$ and $a_j + a_k - 1 < 1$. Thus, after performing redundant blow-ups, the number of redundant points is still finite. $\square$

Proposition 3.6. *The number of surfaces obtained by a sequence of redundant blow-ups from S is finite.*

Proof. Let $\widetilde{p} := \widetilde{E}_j \cap E$ the intersection point. Then we have

$$\mathrm{mult}_{\widetilde{p}}(\widetilde{N}) = a_j + (\mathrm{mult}_p(N) - 1) = \mathrm{mult}_p(N) - (1 - a_j) < \mathrm{mult}_p(N),$$

i.e., the multiplicity strictly decreases after redundant blow-ups.

Suppose that $\mathrm{mult}_{\widetilde{p}} \widetilde{N} \geq 1$. Let $f' \colon \widetilde{S}' \to \widetilde{S}$ be a redundant blow-up at $\widetilde{p}$ with the redundant curve F, and $\widetilde{E}'_j, \widetilde{E}'_k$ and E' be the strict transforms of $\widetilde{E}_j, \widetilde{E}_k$ and E, respectively. Let $\widetilde{p}' := \widetilde{E}'_j \cap F$ and $q := E' \cap F$ be the intersection points.



Let $-K_{\widetilde{S}'} = \widetilde{P}' + \widetilde{N}'$ be the Zariski decomposition. Then we have

$$\begin{aligned}
\mathrm{mult}_{\widetilde{p}}(\widetilde{N}) - \mathrm{mult}_{\widetilde{p}'}(\widetilde{N}') &= \mathrm{mult}_{\widetilde{p}}(\widetilde{N}) - \{\mathrm{mult}_{\widetilde{p}}(\widetilde{N}) - (1 - a_j)\} = 1 - a_j, \\
\mathrm{mult}_{\widetilde{p}}(\widetilde{N}) - \mathrm{mult}_q(\widetilde{N}') &= \mathrm{mult}_{\widetilde{p}}(\widetilde{N}) - \{(\mathrm{mult}_{\widetilde{p}}(\widetilde{N}) - 1) + (\mathrm{mult}_p(N) - 1)\} \\
&= 2 - \mathrm{mult}_p(N) \\
&= 2 - a_j - a_k.
\end{aligned}$$

Note that $\mathrm{mult}_p(N) - \mathrm{mult}_{\widetilde{p}}(\widetilde{N}) = 1 - a_j < 2 - a_j - a_k$, so we obtain

$$\mathrm{mult}_p(N) - \mathrm{mult}_{\widetilde{p}}(\widetilde{N}) \leq \mathrm{mult}_{\widetilde{p}}(\widetilde{N}) - \mathrm{mult}_{\widetilde{p}'}(\widetilde{N}'),$$

and

$$\mathrm{mult}_p(N) - \mathrm{mult}_{\widetilde{p}}(\widetilde{N}) \leq \mathrm{mult}_{\widetilde{p}}(\widetilde{N}) - \mathrm{mult}_q(\widetilde{N}'),$$

i.e., the difference of multiplicities increases after redundant blow-up. Thus, the assertion immediately follows. $\square$

There exist natural numbers M_j and M_k such that

$$\mathrm{mult}_p(N) - M_j(1 - a_j) < 1 \text{ and} \, \mathrm{mult}_p(N) - (M_j - 1)(1 - a_j) \geq 1,$$

and

$$\mathrm{mult}_p(N) - M_k(1 - a_k) < 1 \text{ and} \, \mathrm{mult}_p(N) - (M_k - 1)(1 - a_k) \geq 1.$$

Since $\max\{M_j, M_k\}$ depends only on p, we denote it by $M(p)$.

Corollary 3.7. *The maximal length of sequences of redundant blow-ups from S is equal to*

$$\max_{p \in R}\{M(p)\}$$

where R is the set of all redundant points on S.

3.1.3. *$\bar{S}$ contains at worst rational singularities.*

Lemma 3.8. *If $\bar{S}$ contains a rational singularity that is not a log terminal singularity, then any surface $\widetilde{S}'$ obtained by a sequence of redundant blow-ups from S contains a curve C contracted by the morphism $\widetilde{S}' \to \bar{S}$ such that every point in C is a redundant point. In particular, $\widetilde{S}'$ has infinitely many redundant points.*

Proof. By Remark 3.1, we can choose an integer k with $1 \leq k \leq l$ such that $a_k \geq 1$, so every point in E_k is a redundant point. After a redundant blow-up $\widetilde{S} \to S$ at a point p in E_k, every point in the proper transform of E_k is again a redundant point in $\widetilde{S}$. In this way, $\widetilde{S}'$ contains a curve on which every point is a redundant point. In other words, no matter how many times we perform redundant blow-ups, the resulting surface will contain infinitely many redundant points. □

Proof of Theorem 1.6. By Lemma 3.5 and 3.8, the theorem holds. □

3.2. **Calculation of discrepancies.**

Recall that $\bar{S}$ is a normal projective rational surface with nef anticanonical divisor having at worst rational singularities, and $g\colon S \to \bar{S}$ is the minimal resolution. The aim of this subsection is to prove the following theorem which is nothing but Theorem 1.7 (2).

Theorem 3.9. *S has no redundant point if and only if $\bar{S}$ has at worst canonical singularities or log terminal singularities of type $[\underbrace{2,\dots,2}_{\alpha},3] (\alpha \geq 1)$, $[2,2,3,2]$, $[2,3,2]$, $[2,4]$, or $[n]$ for $n \geq 3$.*

For notation, see Subsubsection 3.2.1. By Lemmas 3.4 and 3.8, we only need to consider the case when $\bar{S}$ has at worst log terminal singularities. Recall that, in this case, $p \in S$ is a redundant point if and only if there are two intersecting irreducible exceptional curves E_j and E_k in the minimal resolution such that the sum of discrepancies $a_j + a_k \geq 1$. Thus, it suffices to consider the problem locally near each singular point s in $\bar{S}$. Throughout this subsection, we assume that $\bar{S}$ contains only one log terminal singular point s.

It is well known that a *dual graph* of a surface rational singularity is always a tree. Each vertex is a smooth rational curve which is an irreducible component of the exceptional divisor of the minimal resolution and the weight is its self-intersection number. Two vertices are connected by an edge if they intersect each other.

In the case of characteristic zero, Brieskorn completely classified finite subgroups of $GL(2,k)$ without quasi-reflections, i.e., he classified all the dual graphs of quotient singularities of surfaces ([Br]). It turns out that the complete list of the dual graphs of the log terminal surface singularities remains the same in arbitrary characteristic (see [A]). For reader's convenience, we will give a detailed description of all possible types ($A_{q,q_1}, D_{q,q_1}, T_m, O_m$ and I_m) of dual graphs and discrepancies of log terminal singularities in following subsubsections.

3.2.1. *A_{q,q_1}-type.*

The dual graph of a log terminal singularity $s \in \bar{S}$ of type A_{q,q_1} is



where each $n_i \geq 2$ is an integer for all i. This singularity can be characterized by the so-called *Hirzebruch-Jung continued fraction*

$$\frac{q}{q_1} = [n_1, n_2, ..., n_l] = n_1 - \cfrac{1}{n_2 - \cfrac{1}{\ddots - \cfrac{1}{n_l}}}.$$

The intersection matrix is

$$M(-n_1, \dots, -n_l) := \begin{pmatrix} -n_1 & 1 & 0 & \cdots & \cdots & 0 \\ 1 & -n_2 & 1 & \cdots & \cdots & 0 \\ 0 & 1 & -n_3 & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & -n_{l-1} & 1 \\ 0 & 0 & 0 & \cdots & 1 & -n_l \end{pmatrix}.$$

For simplicity, we use the notation $[n_1, \dots, n_l]$ to refer a log terminal singularity of A_{q,q_1}-type. Note that $[n_1, \dots, n_l]$ and $[n_l, \dots, n_1]$ denote the same singularity.

We will use the following notation for convenience (cf. [HwK2]).

(1) $q_{b_1,b_2,\dots,b_m} := |\det(M')|$ where M' is the $(l-m) \times (l-m)$ matrix obtained by deleting $-n_{b_1}, -n_{b_2}, \dots, -n_{b_m}$ from $M(-n_1, \dots, -n_l)$. For convenience, we also define $q_{1,\dots,l} = |\det(M(\emptyset))| = 1$.
(2) $u_s := q_{s,\dots,l} = |\det(M(-n_1, \dots, -n_{s-1}))|$ $(2 \le s \le l)$, $u_0 = 0, u_1 = 1$.
(3) $v_s := q_{1,\dots,s} = |\det(M(-n_{s+1}, \dots, -n_l))|$ $(1 \le s \le l-1)$, $v_l = 1, v_{l+1} = 0$.
(4) $q = |\det(M(-n_1, \dots, -n_l))| = u_{l+1} = v_0$.
(5) $|[n_1, \dots, n_l]| := |\det(M(-n_1, \dots, -n_l))|$.

It follows that $q_1 = |\det(M(-n_2, \dots, -n_l)| = v_1$. The following properties of Hirzebruch-Jung continued fractions will be used.

Lemma 3.10. *For $1 \le i \le l$, we have the followings.*

(1) $a_i = 1 - \frac{u_i + v_i}{q}$.
(2) $u_{i+1} = n_i u_i - u_{i-1}$, $v_{i-1} = n_i v_i - v_{i+1}$.
(3) $v_i u_{i+1} - v_{i+1} u_i = v_{i-1} u_i - v_i u_{i-1} = q$.
(4) $|[n_1, \dots, n_{i-1}, n_i + 1, n_{i+1}, \dots, n_l]| = v_i u_i + |[n_1, n_2, \dots, n_l]| > q$.
(5) *If $l \ge 2$ and $n_1 \ge 3$, then $v_1 + v_2 < q$.*

Proof. (1)-(4) are well-known facts. For (1) see [HwK1, Lemma 2.2], and for (2)-(4) see [HwK2, Lemma 2.4]. It is straightforward to check (5) as follow:

$q = v_1 u_2 - v_2 u_1 = v_1 \cdot n_1 - v_2 = (n_1 - 1) v_1 + (v_1 - v_2) > 2 v_1 > v_1 + v_2$. □

Now we give a criterion for checking $a_k + a_{k+1} \ge 1$ for some k. Recall that the intersection point of E_K and E_{k+1} is a redundant point in S.

Lemma 3.11. *The followings are equivalent.*

(1) $a_k + a_{k+1} \ge 1$ *for some* $1 \le k \le l-1$.
(2) $q \ge u_k + u_{k+1} + v_k + v_{k+1}$ *for some* $1 \le k \le l-1$.

Proof. By Lemma 3.10 (1), we have

$$a_k + a_{k+1} = \left(1 - \frac{u_k + v_k}{q}\right) + \left(1 - \frac{u_{k+1} + v_{k+1}}{q}\right) = 2 - \frac{u_k + u_{k+1} + v_k + v_{k+1}}{q},$$

so the equivalence follows. □

The singularity $[\underbrace{2,\dots,2}_{l}]$ is a canonical singularity of type A_l, so $q = l+1$ and $a_i = 0$ for all $1 \le i \le l$. On the other hand, the minimal resolution S of the singularity $[\underbrace{2,\dots,2}_{l-1},3]$ does not contain any redundant point. Indeed, $q = 2l+1$ and $u_i = i$, $v_i = 2l-2i+1$ for $1 \le i \le l$ by using Lemma 3.10. Then, for $1 \le i \le l-1$

$$u_i + u_{i+1} + v_i + v_{i+1} = 4l - 2i + 1 > 2l + 1 = q,$$

so by Lemma 3.11, there is no redundant point on S.

Proposition 3.12. *Let $s \in \bar{S}$ be a log terminal singularity of type A_{q,q_1}. Then S does not contain a redundant point if and only if s is of type $[\underbrace{2,\dots,2}_{\alpha}](\alpha \ge 1)$, $[\underbrace{2,\dots,2}_{\alpha},3](\alpha \ge 1)$, $[2,2,3,2]$, $[2,3,2]$, $[2,4]$, or $[n]$ for $n \ge 3$.*

To prove the proposition, we need two more lemmas.

Lemma 3.13. *The followings hold.*

1. *If s is the singularity $[\underbrace{2,\dots,2}_{\alpha},3,\underbrace{2,\dots,2}_{\beta}]$ $(\alpha \ge \beta \ge 1)$, then $q = \alpha\beta + 2\alpha + 2\beta + 3$ and the intersection point of E_α and $E_{\alpha+1}$ is a redundant point in S except for $[2,2,3,2]$ and $[2,3,2]$.*
2. *If s is the singularity $[\underbrace{2,\dots,2}_{\alpha},3,\underbrace{2,\dots,2}_{\beta},3,\underbrace{2,\dots,2}_{\gamma}]$ $(\alpha \ge 0,\ \beta \ge 1$ and $\gamma \ge 0)$, then the intersection point of $E_{\alpha+1}$ and $E_{\alpha+2}$ is a redundant point in S.*
3. *If s is the singularity $[\underbrace{2,\dots,2}_{\alpha},4,\underbrace{2,\dots,2}_{\beta}]$ $(\alpha \ge \beta \ge 0)$, then the intersection point of E_α and $E_{\alpha+1}$ is a redundant point in S except for $[2,4]$ and $[4]$.*

Proof. The strategy is as follows. First, we compute $u_\alpha, u_{\alpha+1}, u_{\alpha+2}, v_\alpha, v_{\alpha+1}, v_{\alpha+2}$ and q using Lemma 3.10. Second, we determine whether

$$q \ge u_\alpha + u_{\alpha+1} + v_\alpha + v_{\alpha+1} \quad \text{or} \quad q \ge u_{\alpha+1} + u_{\alpha+2} + v_{\alpha+1} + v_{\alpha+2}$$

holds or not. Finally, by Lemma 3.11, we find a redundant point.

(1) We have

$$u_\alpha = |[\underbrace{2,\dots,2}_{\alpha-1}]| = \alpha, v_\alpha = |[3,\underbrace{2,\dots,2}_{\beta}]| = 2\beta+3$$
$$u_{\alpha+1} = |[\underbrace{2,\dots,2}_{\alpha}]| = \alpha+1, v_{\alpha+1} = |[\underbrace{2,\dots,2}_{\beta}]| = \beta+1.$$

Then $u_\alpha + u_{\alpha+1} + v_\alpha + v_{\alpha+1} = 2\alpha + 3\beta + 5$. Using Lemma 3.10 (3), we obtain

$$q = v_\alpha u_{\alpha+1} - v_{\alpha+1} u_\alpha = (\alpha+1)(2\beta+3) - (\beta+1)\alpha = \alpha\beta + 2\alpha + 2\beta + 3.$$

Then we have $q - (u_\alpha + u_{\alpha+1} + v_\alpha + v_{\alpha+1}) = \alpha\beta - \beta - 2 = (\alpha-1)\beta - 2$. If $\alpha\beta - \beta - 2 < 0$, then $(\alpha,\beta) = (1,1)$ or $(2,1)$. These cases are exactly $[2,2,3,2]$ and $[2,3,2]$, and we can easily check that there is no redundant point on S (see Table 1). Otherwise, we have $q \ge (u_\alpha + u_{\alpha+1} + v_\alpha + v_{\alpha+1})$.

(2) We have

$$u_{\alpha+1} = |[\underbrace{2,\dots,2}_{\alpha}]| = \alpha+1, v_{\alpha+1} = |[\underbrace{2,\dots,2}_{\beta},3,\underbrace{2,\dots,2}_{\gamma}]| = \beta\gamma+2\beta+2\gamma+3$$

$$u_{\alpha+2} = |[\underbrace{2,\dots,2}_{\alpha},3]| = 2\alpha+3, v_{\alpha+2} = |[\underbrace{2,\dots,2}_{\beta-1},3,\underbrace{2,\dots,2}_{\gamma}]| = \beta\gamma+2\beta+\gamma+1.$$

Then $u_{\alpha+1}+u_{\alpha+2}+v_{\alpha+1}+v_{\alpha+2} = 2\beta\gamma+3\alpha+4\beta+3\gamma+8$. Using Lemma 3.10 (4), we obtain

$$\begin{aligned} q &= |[\underbrace{2,\dots,2}_{\alpha+\beta+1},3,\underbrace{2\dots,2}_{\gamma}]| + u_{\alpha+1}v_{\alpha+1} \\ &= (\alpha+\beta+1)\gamma+2(\alpha+\beta+1)+2\gamma+3+(\alpha+1)(\beta\gamma+2\beta+2\gamma+3) \\ &= \alpha\beta\gamma+2\alpha\beta+3\alpha\gamma+2\beta\gamma+5\alpha+4\beta+5\gamma+8. \end{aligned}$$

It immediately follows that $q \geq u_{\alpha+1}+u_{\alpha+2}+v_{\alpha+1}+v_{\alpha+2}$.

(3) We have

$$u_{\alpha} = |[\underbrace{2,\dots,2}_{\alpha-1}]| = \alpha, v_{\alpha} = |[4,\underbrace{2,\dots,2}_{\beta}]| = 3\beta+4$$

$$u_{\alpha+1} = |[\underbrace{2,\dots,2}_{\alpha}]| = \alpha+1, v_{\alpha+1} = |[\underbrace{2,\dots,2}_{\beta}]| = \beta+1.$$

Then $u_{\alpha}+u_{\alpha+1}+v_{\alpha}+v_{\alpha+1} = 2\alpha+4\beta+6$. Using Lemma 3.10 (4), we obtain

$$q = |[\underbrace{2,\dots,2}_{\alpha},3,\underbrace{2,\dots,2}_{\beta}]| + u_{\alpha+1}v_{\alpha+1} = 2\alpha\beta+3\alpha+3\beta+4.$$

Then we have $q-(u_{\alpha}+u_{\alpha+1}+v_{\alpha}+v_{\alpha+1}) = 2\alpha\beta+\alpha-\beta-2$. If $2\alpha\beta+\alpha-\beta-2<0$, then $(\alpha,\beta)=(0,0),(1,0)$. These cases are exactly $[2,4]$ and $[4]$, and we can easily check that there is no redundant point on S (see Table 1). Otherwise, we have $q \geq (u_{\alpha}+u_{\alpha+1}+v_{\alpha}+v_{\alpha+1})$. □

Table 1.

singularities	$[2,2,3,2]$	$[2,3,2]$	$[2,4]$	$[n]$
discrepancies	$\left(\frac{2}{11},\frac{4}{11},\frac{6}{11},\frac{3}{11}\right)$	$\left(\frac{1}{4},\frac{1}{2},\frac{1}{4}\right)$	$\left(\frac{2}{7},\frac{4}{7}\right)$	$\left(\frac{n-2}{n}\right)$ $(n \geq 2)$

Lemma 3.14. *Suppose that for* $[n_1,\dots,n_j,\dots,n_k,\dots n_l]$, *there are integers* j *and* k *with* $1 \leq j \leq k \leq l-1$ *such that* $n_j \geq 3$ *and* $u_j+u_{j+1}+v_j+v_{j+1} \leq q$. *Let*

$$[n'_1,\dots,n'_j,\dots,n'_{k-1},n'_k,n'_{k+1},\dots,n'_l] := [n_1,\dots,n_j,\dots,n_{k-1},n_k+1,n_{k+1},\dots,n_l].$$

Then we have $u'_j+u'_{j+1}+v'_j+v'_{j+1} \leq q'$.

Proof. By Lemma 3.10, $q' = q+u_kv_k$. It suffices to show that $u'_j+u'_{j+1}+v'_j+v'_{j+1} \leq q+u_kv_k$. In order to calculate u'_j, u'_{j+1}, v'_j and v'_{j+1}, we divide it into three cases.

Case 1: $j+2 \leq k$.

We have $u'_j = u_j$ and $u'_{j+1} = u_{j+1}$. Moreover, we also have

$$v'_j = |[n_{j+1},\dots,n_{k-1},n_k+1,n_{k+1},\dots,n_l]| = v_j + v_k|[n_{j+1},\dots,n_{k-1}]| \text{ and}$$
$$v'_{j+1} = |[n_{j+2},\dots,n_{k-1},n_k+1,n_{k+1},\dots,n_l]| = v_{j+1} + v_k|[n_{j+2},\dots,n_{k-1}]|$$

Then $u'_j + u'_{j+1} + v'_j + v'_{j+1} \leq q'$ is equivalent to

$$\begin{aligned}&u_j + u_{j+1} + v_j + v_{j+1} + v_k(|[n_{j+1}, \dots, n_{k-1}]| + |[n_{j+2}, \dots, n_{k-1}]|)\\ &\leq q + u_k v_k = q + v_k|[n_j, \dots, n_{k-1}]|.\end{aligned}$$

The above inequality always holds by the assumption and Lemma 3.10 (5).

Case 2: $j + 1 = k$.

We have $u'_j = u_j$ and $u'_{j+1} = u_{j+1}$. Moreover, we also have

$$v'_j = |[n_{j+1} + 1, n_{j+2}, \dots, n_l]| = v_j + v_{j+1} \text{ and } v'_{j+1} = |[n_{j+2}, \dots, n_l]| = v_{j+1}.$$

Since $v_{j+1} \leq u_{j+1}v_{j+1}$, we obtain $u'_j + u'_{j+1} + v'_j + v'_{j+1} \leq q + u_{j+1}v_{j+1}$.

Case 3: $j = k$.

We have $u'_j = u_j$ and $u'_{j+1} = |[n_1, \dots, n_j + 1]| = u_{j+1} + u_j$. Moreover, we also have $v'_j = v_j$ and $v'_{j+1} = v_{j+1}$. As in Case 2, we obtain $u'_j + u'_{j+1} + v'_j + v'_{j+1} \leq q + u_{j+1}v_{j+1}$. □

Proof of Proposition 3.12. First, we introduce notations for simplicity. Fix a natural number l. For integers $n_1, \dots, n_l \geq 2$ and $n'_1, \dots, n'_l \geq 2$, we write $[n_1, \dots, n_l] < [n'_1, \dots, n'_l]$ if and only if $n_i \leq n'_i$ for all $1 \leq i \leq l$ and $n_j < n'_j$ for some $1 \leq j \leq l$. Let $\bar{S}$ (resp. $\bar{S}'$) be a surface with the singularity $[n_1, \dots, n_l]$ (resp. $[n'_1, \dots, n'_l]$), and S (resp. S') be the minimal resolution of $\bar{S}$ (resp., $\bar{S}'$). By Lemma 3.14, if S has a redundant point and $[n_1, \dots, n_l] < [n'_1, \dots, n'_l]$, then S' has a redundant point.

Suppose that $s \in \bar{S}$ is the singularity $[n_1, \dots, n_l]$ other than $[\underbrace{2, \dots, 2}_{\alpha}](\alpha \geq 1)$, $[\underbrace{2, \dots, 2}_{\alpha}, 3](\alpha \geq 1)$, $[2, 2, 3, 2]$, $[2, 3, 2]$, $[2, 4]$, or $[n]$ for $n \geq 3$ in which cases S has no redundant point. Then $[n_1, \dots, n_l]$ is bigger than or equal to $[\underbrace{2, \dots, 2}_{\alpha}, 3, \underbrace{2, \dots, 2}_{\beta}](\alpha \geq \beta \geq 1)$ (not equal to $[2, 2, 3, 2]$ and $[2, 3, 2]$), $[\underbrace{2, \dots, 2}_{\alpha}, 3, \underbrace{2, \dots, 2}_{\beta}, 3, \underbrace{2, \dots, 2}_{\gamma}]$ $(\alpha \geq 0, \beta \geq 1$ and $\gamma \geq 0)$, $[\underbrace{2, \dots, 2}_{\alpha}, 4, \underbrace{2, \dots, 2}_{\beta}]$ $(\alpha \geq \beta \geq 0)$ (not equal to $[2, 4]$ and $[4]$), $[2, 2, 3, 3]$, $[2, 3, 3, 2]$, $[2, 3, 3]$, $[3, 3]$ or $[2, 5]$ in which cases S has a redundant point thanks to Lemma 3.13 and Table 2. Thus, by Lemma 3.14, $[n_1, \dots, n_l]$ has a redundant point. □

TABLE 2.

singularities	$[2, 2, 3, 3]$	$[2, 3, 3, 2]$	$[2, 3, 3]$	$[3, 3]$	$[2, 5]$
discrepancies	$\left(\frac{2}{9}, \frac{4}{9}, \frac{6}{9}, \frac{5}{9}\right)$	$\left(\frac{1}{3}, \frac{2}{3}, \frac{2}{3}, \frac{1}{3}\right)$	$\left(\frac{4}{13}, \frac{8}{13}, \frac{7}{13}\right)$	$\left(\frac{1}{2}, \frac{1}{2}\right)$	$\left(\frac{1}{3}, \frac{2}{3}\right)$

3.2.2. *D_{q,q_1}-type.* The dual graph of a log terminal singularity $s \in \bar{S}$ of D_{q,q_1}-type is



where $\frac{q}{q_1} = [n_1, \ldots, n_l]$, and $b \geq 2$ and $n_i \leq 2$ are integers for all $1 \leq i \leq l$. The matrix equation for each discrepancy a_i is given by

$$\begin{pmatrix} -2 & 0 & 1 & 0 & 0 & 0 & \cdots & 0 \\ 0 & -2 & 1 & 0 & 0 & 0 & \cdots & 0 \\ 1 & 1 & -b & 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1 & -n_1 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 1 & -n_2 & 1 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 1 & -n_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & 1 & -n_l \end{pmatrix} \begin{pmatrix} a_{l+1} \\ a_{l+2} \\ a_0 \\ a_1 \\ a_2 \\ a_3 \\ \vdots \\ a_l \end{pmatrix} = - \begin{pmatrix} 0 \\ 0 \\ b-2 \\ n_1 - 2 \\ n_2 - 2 \\ n_3 - 2 \\ \vdots \\ n_l - 2 \end{pmatrix}.$$

Lemma 3.15 (Lemma 3.7 of [HwK1]). *We have the followings.*

(1) $a_0 = 1 - \frac{1}{(b-1)q-q_1}$ *and* $a_{l+1} = a_{l+2} = \frac{1}{2}a_0$.
(2) $a_1 = 1 - \frac{b-1}{(b-1)q-q_1}$.
(3) $a_l = 1 - \frac{(b-1)q_l - q_{1,l}}{(b-1)q-q_1}$ *for* $l \geq 2$.

Proposition 3.16. *Let* $s \in \bar{S}$ *be a log terminal singularity of* D_{q,q_1}*-type. Then we have the following four cases.*

(1) *If* $b \geq 3$*, then* $a_0 + a_{l+1} \geq 1$.
(2) *If* $b = 2$ *and* $q \geq q_1 + 3$*, then* $a_0 + a_{l+1} \geq 1$.
(3) *If* $b = 2$ *and* $q = q_1 + 2$*, then* $a_0 = a_1 = \frac{1}{2}$.
(4) *If* $b = 2$ *and* $q = q_1 + 1$*, then* s *is a canonical singularity.*

In particular, the minimal resolution S *of* $\bar{S}$ *always has a redundant point unless* s *is a canonical singularity.*

Proof. Using Lemma 3.15, we get

$$a_0 + a_{l+1} = \frac{3}{2}a_0 = \frac{3}{2}\left(1 - \frac{1}{(b-1)q - q_1}\right) \geq 1$$

which is equivalent to

$$(b-1)q - q_1 \geq 3.$$

If $b \geq 3$, then

$$(b-1)q - q_1 \geq 2q - q_1 > q \geq 2.$$

Thus $(b-1)q - q_1 \geq 3$, that is $a_0 + a_{l+1} \geq 1$, so we obtain (1).

Suppose that $b = 2$. If $q - q_1 \geq 3$, then it is still true that $a_0 + a_{l+1} \geq 1$. This is Case (2).

For Case (3), we assume that $b = 2$ and $q = q_1 + 2$. The condition $q = q_1 + 2$ is equivalent to $(n_1 - 1)q_1 = q_{1,2} + 2$. By simple calculation, we conclude that $n_1 = 2$ and $q_1 = q_{1,2} + 2$. By induction,

$$[n_1, \ldots, n_{l-1}, n_l] = [2, \ldots, 2, 3].$$

Using Lemma 3.15, we have $a_0 = a_1 = \frac{1}{2}$, so we are done.

For the last case, assume that $b = 2$ and $q = q_1 + 1$. Similar as Case (3), we obtain

$$[n_1, \dots, n_l] = [2, \dots, 2, 2].$$

Thus y is a canonical singularity of type D_{l+3}. □

3.2.3. *T_m, O_m, and I_m-types.* We use the notation $\langle b; q, q_1; q', q'_1 \rangle$ to refer the following dual graph



where $b \geq 2$ and $n_i \geq 2$ are integers for all $2 \leq i \leq l$, and $\frac{q}{q_1} = [n_2, \dots, n_k]$ and $\frac{q'}{q'_1} = [n_{k+1}, \dots, n_l]$ are the Hirzebruch-Jung continued fractions. Each dual graph of a log terminal singularity $s \in \bar{S}$ of T_m, I_m or O_m-types is one of three items from the top, eight items from the bottom, or the remaining items in Table 3, respectively.

For example, consider $\langle b; 3, 1; 4, 3 \rangle$ which is of O_m-type.



The matrix equation for discrepancies a_i is given by

$$\begin{pmatrix} -b & 1 & 1 & 1 & 0 & 0 \\ 1 & -2 & 0 & 0 & 0 & 0 \\ 1 & 0 & -3 & 0 & 0 & 0 \\ 1 & 0 & 0 & -2 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 & 1 & -2 \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \end{pmatrix} = \begin{pmatrix} 2-b \\ 0 \\ -1 \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$

It is easy to see that the solution $(a_0, a_2, a_2, a_3, a_4, a_5)$ is

$$\left(\frac{12b-20}{12b-19}, \frac{6b-10}{12b-19}, \frac{8b-13}{12b-19}, \frac{9b-15}{12b-19}, \frac{6b-12}{12b-19}, \frac{3b-5}{12b-19} \right).$$

Since $b \geq 2$, we obtain $a_0 + a_1 \geq 1$ by the following computation:

$$\frac{12b-20}{12b-19} + \frac{6b-10}{12b-19} = \frac{18b-30}{12b-19} \geq 1$$

is equivalent to

$$6b \geq 11.$$

Similarly, we calculate all discrepancies of 15 cases. For reader's convenience, we list discrepancies $(a_0, a_1, \dots, a_l)$ of all possible cases in Table 3. It is easy to check that $a_0 + a_1 \geq 1$ for all cases provided that $b \geq 2$ and y is not a canonical singularity. Thus, we obtain the following.

Proposition 3.17. *If $s \in \bar{S}$ is a log terminal singularity of one of T_m, O_m, and I_m-types and s is not a canonical singularity, then $a_0 + a_1 \geq 1$.*

Table 3.

singularities	discrepancies $(a_0, a_1, \dots, a_l)$
$\langle b; 3, 1; 3, 1 \rangle$	$\left(\frac{6b-8}{6b-7}, \frac{3b-4}{6b-7}, \frac{4b-5}{6b-7}, \frac{4b-5}{6b-7}\right)$
$\langle b; 3, 1; 3, 2 \rangle$	$\left(\frac{6b-10}{6b-9}, \frac{3b-5}{6b-9}, \frac{12b-19}{18b-27}, \frac{12b-20}{18b-27}, \frac{6b-10)}{18b-27)}\right)$
$\langle b; 3, 2; 3, 2 \rangle$	$\left(\frac{6b-12}{6b-11}, \frac{3b-6}{6b-11}, \frac{4b-8}{6b-11}, \frac{2b-4}{6b-11}, \frac{4b-8}{6b-11}, \frac{2b-4}{6b-11}\right)$
$\langle b; 3, 2; 4, 3 \rangle$	$\left(\frac{12b-24}{12b-23}, \frac{6b-12}{12b-23}, \frac{8b-16}{12b-23}, \frac{4b-8}{12b-23}, \frac{9b-18}{12b-23}, \frac{6b-12}{12b-23}, \frac{3b-6}{12b-23}\right)$
$\langle b; 3, 1; 4, 3 \rangle$	$\left(\frac{12b-20}{12b-19}, \frac{6b-10}{12b-19}, \frac{8b-13}{12b-19}, \frac{9b-15}{12b-19}, \frac{6b-10}{12b-19}, \frac{3b-5}{12b-19}\right)$
$\langle b; 3, 2; 4, 1 \rangle$	$\left(\frac{12b-18}{12b-17}, \frac{6b-9}{12b-17}, \frac{8b-12}{12b-17}, \frac{4b-6}{12b-17}, \frac{9b-13}{12b-17}\right)$
$\langle b; 3, 1; 4, 1 \rangle$	$\left(\frac{12b-14}{12b-13}, \frac{6b-7}{12b-13}, \frac{8b-9}{12b-13}, \frac{9b-10}{12b-13}\right)$
$\langle b; 3, 2; 5, 4 \rangle$	$\left(\frac{30b-60}{30b-59}, \frac{15b-30}{30b-59}, \frac{20b-40}{30b-59}, \frac{10b-20}{30b-59}, \frac{24b-48}{30b-59}, \frac{18b-36}{30b-59}, \frac{12b-24}{30b-59}, \frac{6b-12}{30b-59}\right)$
$\langle b; 3, 2; 5, 3 \rangle$	$\left(\frac{30b-54}{30b-53}, \frac{15b-27}{30b-53}, \frac{20b-36}{30b-53}, \frac{10b-18}{30b-53}, \frac{24b-43}{30b-53}, \frac{18b-32}{30b-53}\right)$
$\langle b; 3, 1; 5, 4 \rangle$	$\left(\frac{30b-50}{30b-49}, \frac{15b-25}{30b-49}, \frac{20b-33}{30b-49}, \frac{24b-40}{30b-49}, \frac{18b-30}{30b-49}, \frac{12b-20}{30b-49}, \frac{6b-10}{30b-49}\right)$
$\langle b; 3, 2; 5, 2 \rangle$	$\left(\frac{30b-48}{30b-47}, \frac{15b-24}{30b-47}, \frac{20b-32}{30b-47}, \frac{10b-16}{30b-47}, \frac{24b-38}{30b-47}, \frac{12b-19}{30b-47}\right)$
$\langle b; 3, 1; 5, 3 \rangle$	$\left(\frac{30b-44}{30b-43}, \frac{15b-22}{30b-43}, \frac{20b-29}{30b-43}, \frac{24b-35}{30b-43}, \frac{18b-26}{30b-43}\right)$
$\langle b; 3, 2; 5, 1 \rangle$	$\left(\frac{30b-42}{30b-41}, \frac{15b-21}{30b-41}, \frac{20b-28}{30b-41}, \frac{10b-14}{30b-41}, \frac{24b-33}{30b-41}\right)$
$\langle b; 3, 1; 5, 2 \rangle$	$\left(\frac{30b-38}{30b-37}, \frac{15b-19}{30b-37}, \frac{20b-25}{30b-37}, \frac{24b-30}{30b-37}, \frac{12b-15}{30b-37}\right)$
$\langle b; 3, 1; 5, 1 \rangle$	$\left(\frac{30b-32}{30b-31}, \frac{15b-16}{30b-31}, \frac{20b-21}{30b-31}, \frac{24b-25}{30b-31}\right)$

Proof of Theorem 3.9. It is a consequence of Lemmas 3.8, 3.4, Propositions 3.12, 3.16, and 3.17. ☐

Proof of Theorem 1.7. (1) follows from Lemma 3.8 and Proposition 3.6, and (2) follows from Theorem 3.9. ☐

4. Rational surfaces with big anticanonical divisor

Question 4.1 (see Remark 3 of [TVAV])**.** *Is every big anticanonical rational surface the minimal resolution of a del Pezzo surface containing at worst rational singularities?*

In this section, we answer Question 4.1 by constructing big anticanonical rational surfaces that are not minimal resolutions of del Pezzo surfaces with rational singularities, and discuss the characterization of weak del Pezzo surfaces.

4.1. **Proof of Theorem 1.9.** Recall that the anticanonical morphism from a big anticanonical rational surface $\widetilde{S}$ factors through the minimal resolution S of a del Pezzo surface $\bar{S}$ with rational singularities, followed by a sequence of redundant blow-ups (see Theorem 2.12). Thus, Question 4.1 reduces to the problem of finding redundant points on S, since in this case the minimal resolution coincides with the anticanonical morphism.

First, we give an explicit construction of big anticanonical rational surfaces whose anticanonial models have at worst log terminal singularities.

Example 4.2. Let $\pi\colon S(m,n) \to \mathbb{P}^2$ be the blow-up of $\mathbb{P}^2$ at $p^1_1, \dots p^1_m$ on a line $\bar{l}_1$ and $p^2_1, \dots p^2_n$ on on a different line $\bar{l}_2$ for $m \geq n \geq 4$. Assume that all points are distinct and away from the intersection point of $\bar{l}_1$ and $\bar{l}_2$. Let l_i be the strict transform of $\bar{l}_i$, E^i_j be the exceptional divisors of p^i_j, and l be the pull-back of a general line in $\mathbb{P}^2$. Then the anticanonical divisor of $S(m,n)$ is given by

$$-K_{S(m,n)} = \pi^*(-K_{\mathbb{P}^2}) - \sum_{i,j} E^i_j = l + l_1 + l_2.$$

Let $-K_{S(m,n)} = P + N$ be the Zariski decomposition. Then we have

$$N = \frac{mn-m-2n}{mn-m-n} l_1 + \frac{mn-2m-n}{mn-m-n} l_2.$$

It is easy to see that $X(m,n)$ is a big anticanonical rational surface. Now the intersection point p of l_1 and l_2 is a redundant point because

$$\frac{mn-m-2n}{mn-m-n} + \frac{mn-2m-n}{mn-m-n} \geq 1.$$

Thus, by blowing up at p, we obtain a redundant blow-up $f\colon \widetilde{S}(m,n) \to S(m,n)$. The Picard number of $\widetilde{S}(m,n)$ is $m+n+2$, so it supports Theorem 1.9. In this case, $M_1 > m-2-\frac{2m}{n}$ and $M_2 > n-2-\frac{2n}{m}$, so $M(x)$ can be increased arbitrarily large (see Subsection 3.1 for notation).

In the next example, we construct big anticanonical rational surfaces whose anticanonical models have at least one rational singularity that is not a log terminal singularity.

Example 4.3. Let $h\colon \mathbb{F}_n \to \mathbb{P}^1$ be the Hirzebruch surface with a section $\overline{\sigma}$ of self-intersection $-n \leq -2$. Let k be an integer such that $3 \leq k \leq n+1$ and let $a_1, \dots, a_k$ be positive integer such that $\sum_{j=1}^k \frac{1}{a_j} < k-2$. Choose k distinct fibres $\overline{F_1}, \dots, \overline{F_k}$ of h, and choose a_i distinct points $p^i_1, \dots, p^i_{a_i}$ on $\overline{F_i} \backslash \overline{\sigma}$. Let S be the blow-up of $\mathbb{F}_n$ at p^i_j where $1 \leq i \leq k, 1 \leq j \leq a_i$ and σ be the strict transform of $\overline{\sigma}$, and F_i be the strict transform of $\overline{F_i}$. Denote the pull-back of a general fiber of $\mathbb{F}_n$ by F.

Then, we have the Zariski decomposition of an anticanonical divisor $-K_S = P + N$ as follows:

$$P = \frac{n+2-k}{n-\sum \frac{1}{a_j}} \sigma + (n+2-k)F + \sum_{i=1}^k \frac{n+2-k}{a_i(n-\sum \frac{1}{a_j})} F_i,$$

$$N = \left(2 - \frac{n+2-k}{n - \sum \frac{1}{a_j}}\right)\sigma + \sum_{i=1}^{k}\left(1 - \frac{n+2-k}{a_i(n - \sum \frac{1}{a_j})}\right)F_i.$$

Thus, S is a big anticanonical rational surface, and every point p in σ is a redundant point because

$$\mathrm{mult}_p(N) \geq 2 - \frac{n+2-k}{n - \sum \frac{1}{a_j}} > 1.$$

The construction of S appeared in Section 3 of [TVAV].

Singularities on log del Pezzo surfaces are classified in some cases ([AN], [Na], [OT], [Ko], [KKN], [Sü]). Thus we can find redundant points on the minimal resolution by Theorem 1.6 and Theorem 1.7. In this way, we can find many other examples of the big anticanonical rational surfaces that is not a minimal resolution of a rational surface with rational singularities.

Remark 4.4. Let S be a big anticanonical rational surface, and $f\colon \widetilde{S} \to S$ be a blow-up at $p \in S$. It is possible that, even though f is not a redundant blow-up, $\widetilde{S}$ can be a big anticanonical rational surface. However, in this case, the anticanonical model of $\widetilde{S}$ is different from that of S.

4.2. **Singularities from big anticanonical rational surfaces.** The following proposition is well-known. Here we prove it using the anticanonical morphism.

Proposition 4.5. *Let $\bar{S}$ be a normal projective rational surface and $g\colon S \to \bar{S}$ be its minimal resolution. If S is a big anticanonical rational surface, then $\bar{S}$ has at worst rational singularities.*

Proof. Let $-K_S = P + N$ be the Zariski decomposition. Then the anticanonical morphism h is given by $|mP|$ for some integer $m > 0$. We claim that if g contracts a curve C, then so is h. By the adjunction formula,

$$P.C + N.C = -K_S.C = C^2 - 2p_a(C) + 2 \leq 0,$$

because $C^2 \leq -2$ for the case $p_a(C) = 0$, and $C^2 \leq -1$ for any case. Since $P.C \geq 0$, we have $N.C \leq 0$. If $N.C < 0$, then C is contained in the support of N. If $N.C = 0$, then $P.C = 0$. In any case, C is also contracted by h.

Note that every subgraph of a dual graph of a surface rational singularity is also a dual graph of a different surface rational singularity. Thus, every singular point s in $\bar{S}$ is rational. ☐

4.3. **Characterization of weak del Pezzo surfaces.** A *weak del Pezzo surface* is a normal projective rational surface with rational singularities such that the anticanonical divisor is nef and big. In the proof of the following theorem, we study contractions from big anticanonical rational surfaces with redundant curves.

Theorem 4.6. *The following classes are equivalent:*

1. $\{$*minimal resolutions of weak del Pezzo surfaces*$\}$
2. $\{\widetilde{S} : \widetilde{S}$ *is a big anticanonical rational surface such that any redundant curve E is not contained in the support of $\widetilde{N}$ where $-K_{\widetilde{S}} = \widetilde{P} + \widetilde{N}$ is the Zariski decomposition*$\}$

First, we determine negative rational curves C, not supported in N, with $P.C = 0$. The following lemma can be easily obtained from Lemma 2.3.

Lemma 4.7. *Let $f\colon \widetilde{S} \to S$ be redundant blow-up of a big anticanonical rational surface S at a point p, and E be the redundant curve. Let $-K_{\widetilde{S}} = \widetilde{P} + \widetilde{N}$ and $-K_S = P + N$ be the Zariski decompositions. Then E is not contained in the support of $\widetilde{N}$ if and only if* $\operatorname{mult}_p(N) = 1$.

The following proof is almost same to [Sa, Proof of Theorem 4.3], but it contains somewhat different argument (contracting negative curves after redundant blow-ups), hence we include the whole proof for the convenience of the reader.

Proof of Theorem 4.6. Let $\bar{S}'$ be a weak del Pezzo surface, and $h'\colon \widetilde{S} \to \bar{S}'$ be the minimal resolution. By Lemma 3.2, the Zariski decomposition $-K_{\widetilde{S}} = \widetilde{P} + \widetilde{N}$ is given by $\widetilde{P} = h'^*(-K_{\bar{S}'})$, and every irreducible component of $\widetilde{N}$ is contained in the set of exceptional curves of h'. In particular, $\widetilde{S}$ is a big anticanonical rational surface. Also, $\widetilde{N}$ contains no redundant curves, since h' is the minimal resolution. Thus, $\widetilde{S}$ is a member of the second class in the theorem.

Conversely, let $\widetilde{S}$ be a member of the second class, and $h\colon \widetilde{S} \to \bar{S}$ be the anticanonical morphism to a del Pezzo surface $\bar{S}$ with rational singularities. By Theorem 2.12, we have the following diagram

$$\begin{tikzcd}
\widetilde{S} \arrow[rr, "h"] \arrow[dr, "f"'] & & \bar{S} \\
& S \arrow[ur, "g"'] &
\end{tikzcd}$$

where f is a sequence of redundant blow-ups, and g is the minimal resolution. Denote the Zariski decompositions by $-K_{\widetilde{S}} = \widetilde{P} + \widetilde{N}$ and $-K_S = P + N$. Let $E_1, \ldots, E_l$ be irreducible components of the g-exceptional divisor. Suppose that only $E_1, \ldots, E_k$ appear in the support of N. Then the anticanonical morphism $g\colon S \to \bar{S}$ contracts $E_1, \ldots, E_k$ to non-canonical singularities, and $E_{k+1}, \ldots, E_l$ to canonical singularities.

Let $\widetilde{E}_1 := f_*^{-1}E_1, \ldots, \widetilde{E}_l := f_*^{-1}E_l$ be their strict transforms. By Lemma 2.3 and the assumption, we have $\widetilde{N} = f_*^{-1}N$. By construction, the anticanonical morphism $h\colon \widetilde{S} \to \bar{S}$ contracts $\widetilde{E}_1, \ldots, \widetilde{E}_k$ to non-canonical singularities, and $\widetilde{E}_{k+1}, \ldots, \widetilde{E}_l$ to canonical singularities. Let $A := \{\widetilde{E}_1, \ldots, \widetilde{E}_k\}$ be the support of $\widetilde{N}$. By contracting $\widetilde{E}_1, \ldots, \widetilde{E}_l$, we get a proper birational morphism $h'\colon \widetilde{S} \to \bar{S}'$ to a normal projective surface $\bar{S}'$.

$$\begin{tikzcd}
\widetilde{S} \arrow[r, "f"] \arrow[d, "h'"'] & S \arrow[d, "g"] \\
\bar{S}' & \bar{S}
\end{tikzcd}$$

By construction, h' is the minimal resolution of $\bar{S}'$, and contracts $\widetilde{E}_1, \ldots, \widetilde{E}_k$ to non-canonical rational singularities and $\widetilde{E}_{k+1}, \ldots, \widetilde{E}_l$ to canonical singularities.

Now we have

$$\widetilde{N}.\widetilde{E}_i = -K_{\widetilde{S}}.\widetilde{E}_i = (-K_{\widetilde{S}} + h'^*K_{\bar{S}'}).\widetilde{E}_i$$

for any irreducible curve $\widetilde{E}_i$ in A since $\widetilde{P}.\widetilde{E}_i = P.E_i = 0$. Since the intersection matrix of elements of A is negative definite, and the supports of $\widetilde{N}$ and $-K_{\widetilde{S}}+h'^*K_{\bar{S}'}$

coincide, we obtain $\widetilde{N} = -K_{\widetilde{S}} + h'^*K_{\bar{S}'}$ and $\widetilde{P} = h'^*(-K_{\bar{S}'})$. For an irreducible curve $\bar{C}'$ on $\bar{S}'$, let $\widetilde{C}' := h'^{-1}_*\bar{C}'$ be the strict transform. Then we have

$$-K_{\bar{S}'}.\bar{C}' = h'^*(-K_{\bar{S}'}).h'^*\bar{C}' = \widetilde{P}.\widetilde{C}' \geq 0.$$

Thus $-K_{\bar{S}'}$ is nef. Moreover, $(-K_{\bar{S}'})^2 = \widetilde{P}^2 > 0$, so $-K_{\bar{S}'}$ is nef and big. We remark that for the image $h(\bar{E})$ of a redundant curve E on $\widetilde{S}$, we have $-K_{\bar{S}}.\bar{E} = 0$. ☐

Remark 4.8. In Proof of Theorem 4.6, if we take the birational morphism $h'': \widetilde{S} \to \bar{S}''$ contracting $\widetilde{E}_1, \ldots, \widetilde{E}_k$, then we can prove that $\bar{S}''$ is also a weak del Pezzo surface.

5. Minimal resolutions of rational $\mathbb{Q}$-homology projective planes

In this section, we prove Theorem 1.10, and investigate the relation between Mori dream rational surfaces and minimal resolutions of rational $\mathbb{Q}$-homology projective planes.

Proof of Theorem 1.10. Let $\bar{S}$ be a rational surface with log terminal singularities having Picard number one, and $g: S \to \bar{S}$ be the minimal resolution. We assume that $-K_{\bar{S}}$ is nef, so we have two cases: $-K_{\bar{S}}$ is ample, or $-K_{\bar{S}}$ is numerically trivial. If $-K_{\bar{S}}$ is ample, then S is a big anticanonical rational surface. Thus, the assertion immediately follows from [TVAV, Theorem 1] or [CS, Theorem 3].

Now assume that $-K_{\bar{S}}$ is numerically trivial and $\bar{S}$ does not contain any canonical singularity. By Lemma 3.2, the Zariski decomposition $-K_S = P + N$ is given by

$$P = g^*(-K_{\bar{S}}) = 0 \text{ and } N = \sum_{i=1}^{l} a_i E_i,$$

where each E_i denotes an irreducible component of the g-exceptional divisor. Note that

$$\mathrm{Pic}(S) = g^*(\mathrm{Pic}(\bar{S})) \oplus \bigoplus_{i=1}^{l} \mathbb{Z} \cdot E_i,$$

and by the assumption, we have $0 < a_i < 1$ for all i.

First, we show that the effective cone $\mathrm{Eff}(S)$ is rational polyhedral. By Lemma 2.6, it suffices to show that there are only finitely many (-1)-curves and (-2)-curves on S. Let H be an ample generator of $\mathrm{Pic}(\bar{S})$. We may write every integral curve not in the support of N as

$$C = g^*(bH) + \sum_i b_i E_i.$$

Note that $b \geq 0$. Suppose that C is a (-1)-curve. We denotes the nonnegative integer $\lambda_i = C.E_i$ for each i. Then, we have

$$1 = C.(-K_S) = C.N = \sum_i a_i C.E_i = \sum_i a_i \lambda_i.$$

Since each a_i is a positive rational number, there are only finitely many possibilities for each λ_i. We have

$$C.E_j = g^*(bH).E_j + \sum_i b_i E_i.E_j = \sum_i b_i E_i.E_j,$$

so we obtain the matrix equation

$$\begin{pmatrix} E_1^2 & E_2E_1 & \cdots & E_lE_1 \\ E_2E_1 & E_2^2 & \cdots & E_lE_2 \\ \vdots & \vdots & \ddots & \vdots \\ E_lE_1 & E_lE_2 & \cdots & E_l^2 \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_l \end{pmatrix} = \begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \vdots \\ \lambda_l \end{pmatrix}$$

which has a unique solution $(b_1, b_2, \dots, b_l)$ because the intersection matrix in the left hand side is negative definite. Moreover, we have

$$-1 = C^2 = f^*(bH)^2 + (\sum_i b_iE_i)^2,$$

and hence, $\pi^*(bH)^2$ is determined by b_i, so is b. Thus, there are only finitely many (-1)-curves on S.

Suppose that C is a (-2)-curve. Then, we have

$$-K_S.C = N.C = \sum_i a_iE_i.C = 0,$$

so $E_i.C = 0$ for all i. Thus, we may write $C = g^*(bH)$ for some $b \geq 0$, so $C^2 \geq 0$, which is a contradiction. Hence, every (-2)-curve is contained in the support of N.

Finally, we show that every nef effective $\mathbb{Q}$-divisor M is semiample. For sufficiently small $\epsilon > 0$, the pair $(S, N + \epsilon M)$ is Kawamata log terminal, and $K_S + N + \epsilon M = \epsilon M$ is nef. By the log abundance theorem for surfaces (see [FM] for $\operatorname{char}(k) = 0$ and [Ta] for $\operatorname{char}(k) > 0$), M is semiample. □

Remark 5.1. Since $\bar{S}$ does not contain any canonical singularities, every g-exceptional curve appears in N. It plays an important role in showing that S contains finitely many (-1)-curves. There exists a rational surface with numerically trivial anticanonical divisor containing canonical singularities (see [HwK3, Section 6]). We do not know whether the minimal resolution of such surface is a Mori dream rational surface or not.

Remark 5.2. (S, N) is a *Calabi-Yau pair*, which is a Kawamata log terminal pair such that $K_S + N$ is numerically trivial, in the sense of [To], and the log terminal model of $\bar{S}$.

In the rest of the section, we assume that $k = \mathbb{C}$. Recall that a *rational* $\mathbb{Q}$-*homology projective planes* $\bar{S}$ is, by definition, a complex normal projective rational surface with at worst quotient singularities and the second topological Betti number $b_2(\bar{S}) = 1$. The following is an immediate consequence of Theorem 1.10.

Corollary 5.3. *Let $\bar{S}$ be a rational $\mathbb{Q}$-homology projective plane, and $g\colon S \to \bar{S}$ be its minimal resolution. Assume that $-K_{\bar{S}}$ is nef and $\bar{S}$ does not contain any rational double point. Then S is a Mori dream space.*

If the anticanonical divisor $-K_{\bar{S}}$ of the rational $\mathbb{Q}$-homology projective plane is numerically trivial, then the anticanonical Iitaka dimension $\kappa(-K_S) = 0$ for its minimal resolution S, since $P = 0$ for the Zariski decomposition $-K_S = P + N$. Using Theorem 1.10, we construct examples of Mori dream rational surfaces with anticanonical Iitaka dimension 0.

Example 5.4. In [HwK3], Hwang and Keum constructed rational $\mathbb{Q}$-homology projective planes with numerically trivial anticanonical divisor not containing any rational double points. By Theorem 1.10, minimal resolution of those surfaces are Mori dream rational surfaces with anticanonical Iitaka dimension 0.

The construction in Section 4 of [HwK3] is as follows. Let $\bar{l}_1, \bar{l}_2, \bar{l}_3, \bar{l}_4$ be four general lines on $\mathbb{P}^2$, and choose four intersection points. Let $Z(2,2,2,2) \to \mathbb{P}^2$ be the blow-up at the four intersection points twice each. Let E_1, E_2, E_3, E_4 be the four exceptional curves of the first kind. We may assume that each E_i meets the strict transform l_i of $\bar{l}_i$ for $i = 1,2,3,4$. By blowing-up the four points $E_i \cap l_i$ for $i = 1,2,3,4$ in $Z(2,2,2,2)$, we get $Z(3,3,3,3)$ which is the minimal resolution of a rational $\mathbb{Q}$-homology projective plane $T(3,3,3,3)$ with numerically trivial anticanonical divisor. Moreover, $T(3,3,3,3)$ has two cyclic quotient singularities of the same type $[2,2,3,3,2,2]$.

In Section 5 of [HwK3], the minimal resolution $Z(2)$ of rational $\mathbb{Q}$-homology projective plane $S(2)$ with numerically trivial anticanonical divisor containing a unique cyclic quotient singularity of type $[3,2,2,2,2,2,2,2,2,3]$ is also explicitly constructed.

Remark 5.5. Each Mori dream rational surface in Example 5.4 is obtained by blow-ups of a big anticanonical rational surface at one point. They differ from the Mori dream rational surface $\widetilde{Y}$ with $\kappa(-K_{\widetilde{Y}}) = 0$ in [LT, Section 6], because the Zariski decompositions of anticanonical divisors are different. On the other hand, $\widetilde{Y}$ and $Z(2)$ are Coble rational surfaces in the sense of Dalgachev and Zhang ([DZ]), but $Z(3,3,3,3)$ is not a Coble rational surface.

Mori dream rational surfaces in Example 5.4 have redundant points by Theorem 3.9. The surfaces obtained by redundant blow-ups from these surfaces are still Mori dream spaces, even in the case of $\kappa(-K_S) = 0$ and $\operatorname{mult}_p N = 1$. In fact, we prove more general result.

Proposition 5.6. *Let (S, Δ) be a Calabi-Yau pair in dimension two. Let $f\colon \widetilde{S} \to S$ be the blow-up at a point p. Assume that $(\widetilde{S}, f_*^{-1}\Delta)$ is a Calabi-Yau pair. If S is a Mori dream surface, then so is $\widetilde{S}$.*

Proof. Note that the image of $\operatorname{Aut}(S, \Delta)$ in $GL(N^1(S))$ is finite by [To, Corollary 5.1]. By [To, Corollary 5.1], it is sufficient to show that the image of $\operatorname{Aut}(\widetilde{S}, f_*^{-1}\Delta)$ in $GL(N^1(\widetilde{S}))$ is isomorphic to the image of $\operatorname{Aut}(S, \Delta)$ in $GL(N^1(S))$. Any automorphism $\widetilde{\varphi}$ in $\operatorname{Aut}(\widetilde{X}, f_*^{-1}\Delta)$ induces a birational map $\varphi\colon S \dashrightarrow S$ such that φ is defined on $S\backslash\{p\}$ and fixes $\Delta\backslash\{p\}$. The total transform $\varphi(p)$ of p consists of one point, so φ is a morphism by the Zariski main theorem. □

The technical assumption $k = \mathbb{C}$ is essential, since it is used in the main ingrediant [To, Corollary 5.1]. By redundant blow-ups from rational surfaces in Example 5.4, we get more Mori dream rational surfaces with anticanonical Iitaka dimension 0. Furthermore, we obtain the following (cf. Theorem 1.9).

Corollary 5.7. *There exists a Mori dream rational surface $\widetilde{Z}$ with $\kappa(-K_{\widetilde{Z}})= 0$ such that $(\widetilde{Z}, -K_{\widetilde{Z}})$ is a Calabi-Yau pair but not a minimal resolution of a normal projective rational surface S with rational singularities such that $-K_S$ is nef.*

Proof. Consider the rational surface $Z := Z(3,3,3,3)$ in Example 5.4. Let $-K_Z = P + N$ be the Zariski decomposition. By simple calculation, one can easily see that the intersection p of two (-3)-curves in Z is a redundant point and $\operatorname{mult}_p(N) =$

$\frac{3}{2}$. Let $f\colon \widetilde{Z} \to Z$ be the redundant blow-up at p, and $-K_{\widetilde{Z}} = \widetilde{P} + \widetilde{N}$ be the Zariski decomposition. By Theorem 1.1, $\widetilde{Z}$ is a Mori dream rational surface with $\kappa(-K_{\widetilde{Z}}) = 0$, and $(\widetilde{Z}, -K_{\widetilde{Z}})$ is a Calabi-Yau pair. Now the exceptional curve E of f is contained in the support of $\widetilde{N}$ by Lemma 2.3.

Suppose that $\widetilde{Z}$ is the minimal resolution of a normal projective rational surface S with rational singularities such that $-K_S$ is nef. Then by Lemma 3.2, $\widetilde{N}$ does not contain (-1)-curve, which is a contradiction, since $\widetilde{N}$ contains a (-1)-curve E. $\square$

6. Examples of Mori dream rational surfaces with $\kappa(-K) \leq 0$

In this section, we prove Theorem 1.8 by constructing Mori dream rational surfaces with effective k^*-action. For generalities on k^*-surfaces, see [OW] and [HS].

Example 6.1. We have an effective k^*-action on $\mathbb{P}^2 = \mathrm{Proj}(k[x,y,z])$ given by $t \cdot (x,y,z) = (tx,ty,z)$ for $t \in k^*$. Consider k^*-invariant lines $l_1 = V(x), l_2 = V(y), l_3 = V(x+y)$ and $L_1 = V(x-y), L_2 = V(z)$ in $\mathbb{P}^2$. Let $\phi\colon S \to \mathbb{P}^2$ be the blow-up at the base points of the cubic pencil determined by $xy(x+y)$ and $(x-y)z^2$ to make an elliptic fibration $\pi\colon S \to \mathbb{P}^1$ with two singular fibres of the same type $\widetilde{D}_4$. For simplicity, we use the same notations for strict transforms of k^*-invariant lines. In the following picture, solid lines are (-2)-curves and dashed lines are (-1)-curves on S.



If we blow-up at a k^*-fixed point, then k^*-action uniquely extends ([OW, Lemma 1.8]). Thus, k^*-action on $\mathbb{P}^2$ extends to S. Recall that smooth projective rational surfaces with effective k^*-action are Mori dream surfaces ([Kn] and [HS, Theorem 1.3]). Thus, S is a Mori dream rational surface with $\kappa(-K_S) = 1$. There is a section E on the elliptic fibration $\pi\colon S \to \mathbb{P}^1$ meeting two singular fibres at p and q, respectively. Since any negative self-intersection curve is k^*-invariant ([OW, Proposition 1.9]), p and q are k^*-invariant.

Let $f(0)\colon S_p(0) \to X$ be the blow-up at p. For simplicity, we also use the same notation for strict transforms of irreducible components of singular fibres of $\pi\colon S \to \mathbb{P}^1$. Note that $-K_S = 2L_2 + L_1 + E_1 + E_2 + E_3$. The Zariski decomposition $-K_{S_p(0)} = P + N$ is given by $P = 0$ and $N = 2L_2 + L_1 + E_1 + E_2 + E_3$, so $\kappa(-K_{S_p(0)}) = 0$. Since the intersection points of negative self-intersection curves are k^*-invariant points, there is a k^*-invariant point on some component of N. Let

$f(1)\colon S_p(1) \to S_p(0)$ be the blow-up at a k^*-invariant point on N. Then $S_p(1)$ has k^*-action, and $f(1)$ is a redundant blow-up. We can repeat this construction $f(n)\colon S_p(n) \to S_p(n-1)$ for each $n \geq 2$. Note that $S_p(n)$ is a Mori dream rational surface with $\kappa(-K_{S_p(n)}) = 0$ of Picard number $n+11$ for each $n \geq 0$.

For each $n \geq 0$, let $g(n)\colon \widetilde{S}_q(n) \to S_p(n)$ be the blow-up at the k^*-fixed point q. Note that there is a birational surjective morphism $h(n)\colon \widetilde{S}_q(n) \to \widetilde{S}_q(n-1)$. We claim that $\kappa(-K_{\widetilde{S}_q(n)}) = -\infty$. Indeed, if $\kappa(-K_{\widetilde{S}_q(n)}) = 0$, then $g(n)$ is a redundant blow-up by Theorem 1.4. But for the Zariski decomposition $-K_{S_p(n)} = P(n)+N(n)$, the point q is not in the support of $N(n)$, so $\mathrm{mult}_q(N(n)) = 0$, because $N(n)$ is in the fixed part of $|-K_{S_p(n)}|$ by [Sa, Lemma 2.4]. This is a contradiction. Thus, $\widetilde{S}_q(n)$ is a Mori dream rational surface with $\kappa(-K_{\widetilde{S}_q(n)}) = -\infty$ of Picard number $n+12$ for each $n \geq 0$.

$$\begin{tikzcd}
\cdots \arrow[r, "h(2)"] & \widetilde{S}_q(1) \arrow[r, "h(1)"] \arrow[d, "g(1)"] & \widetilde{S}_q(0) \arrow[d, "g(0)"] & & \\
\cdots \arrow[r, "f(2)"] & S_p(1) \arrow[r, "f(1)"] & S_p(0) \arrow[r, "f(0)"] & S \arrow[r, "\phi"] \arrow[d, "\pi"'] & \mathbb{P}^2 \\
& & & \mathbb{P}^1 &
\end{tikzcd}$$

Remark 6.2. We assume that $k = \mathbb{C}$. If we contract E on S, then we get the minimal resolution of a degree one del Pezzo $\mathbb{C}^*$-surface of Picard number one with two canonical singularities of the same type D_4. Del Pezzo $\mathbb{C}^*$-surfaces of Picard number one with canonical singularities are completely classified (see [Sü, Theorem 4.6] and [HS, Theorem 5.6]). If the degree is one, then there are exactly four possible singularity types $2D_4, E_6 + A_2, E_7 + A_1$ or E_8. On the other hand, S is an extremal rational elliptic surface X_{11} in the sense of Miranda and Persson (see [MP, Theorem 4.1]). In general, for each minimal resolution Y of degree one del Pezzo $\mathbb{C}^*$-surfaces of Picard number one with canonical singularities, there exists a $\mathbb{C}^*$-fixed point y on Y such that X is an extremal rational elliptic surface where $X \to Y$ is the blow-up at y. Note that the four extremal rational elliptic surfaces X_{22}, X_{33}, X_{44}, and X_{11} in [MP, Theorem 4.1] have an effective $\mathbb{C}^*$-action. Thus, we can do the similar construction for these elliptic surfaces as in Example 6.1.

Remark 6.3. There exists a redundant point p' on $S_p(n)$ for $n \geq 0$ such that $\mathrm{mult}_{p'}(N(n)) = 1$. If p' is not a k^*-invariant point, then we do not know whether the surface obtained by blowing up at the point p' is a Mori dream space.

References

[A] V. Alexeev, *"Classification of log canonical surface singularities: arithmetical proof"* in *Flips and abundance for algebraic threefolds (Salt lake city, 1991)*, Astérisque **211** (1992), Soc. Math. France, Paris, 47-58.

[AGL] M. Artebani, A. Garbagnati, and A. Laface, *Cox rings of extremal rational elliptic surfaces*, preprint, arXiv:1302.4361.

[AL] M. Artebani and A. Laface, *Cox rings of surfaces and the anticanonical Iitaka dimension*, Adv. Math. **226** (2011), no.6, 5252-5267.

[AL′] M. Artebani and A. Laface, *Cox rings of surfaces and the anticanonical Iitaka dimension*, revised version.

[AN] V. Alexeev and V.V. Nikulin, *Del Pezzo and K3 surfaces*, MSJ Memoirs **15** (2006), Mathematical Society of Japan, Tokyo.

[Ba] L. Bădescu, *Algebraic surfaces*, Universitext. Springer-Verlag, New York, 2001.

[Br] E. Brieskorn, *Rationale Singularitäten komplexer Flächen*, Invent. Math. **4** (1967/1968), 336-358.

[CS] D. Chen and C. Schnell, *Surfaces with big anticanonical class*, preprint (2008).

[DZ] I.V. Dolgachev and D.-Q. Zhang, *Coble rational surfaces*, Amer. J. Math. **123** (2001), no.1, 79-114.

[EH] D. Eisenbud and J. Harris, *"On varieties of minimal degree (a centennial account)"* in *Algebraic geometry, Bowdoin (Brunswick, Maine, 1985)*, Proc. Sympos. Pure Math. **46** (1987), Part 1, Amer. Math. Soc., Providence, RI, 3-13.

[EKW] E.J. Elizondo, K. Kurano, and K. Watanabe, *The total coordinate ring of a normal projective variety*, J. Algebra **276** (2004), no.2, 625-637.

[FM] L.-Y. Fong and J. McKernan, *"Log abundance for surfaces"* in *Flips and abundance for algebraic threefolds (Salt lake city, 1991)*, Astérisque **211** (1992), Soc. Math. France, Paris, 127-137.

[GM] C. Galindo and F. Monserrat, *The total coordinate ring of a smooth projective surface*, J. Algebra **284** (2005), no.1, 91-101.

[H] J. Hausen, *Cox rings and combinatorics II*, Moscow Math. J. **8** (2008), no.4, 711-757.

[HS] J. Hausen and H. Süß, *The Cox ring of an algebraic variety with torus action*, Adv. Math. **225** (2010), no.2, 977-1012.

[HuK] Y. Hu and S. Keel, *Mori dream spaces and GIT*, Michigan Math. J. **48** (2000), 331-348.

[HwK1] D. Hwang and J. Keum, *The maximum number of singular points on rational homology projective planes*, J. Algebraic Geom. **20** (2011), no.3, 495-523.

[HwK2] D. Hwang and J. Keum, *Algebraic Montgomery-Yang problem: the non-rational surface case*, to appear in Michigan Math. J.

[HwK3] D. Hwang and J. Keum, *Construction of singular rational surfaces of Picard number one with ample canonical divisor*, Proc. Amer. Math. Soc. **140** (2012), no.6, 1865-1879.

[KKN] A. Kasprzyk, M. Kreuzer, and B. Nill, *On the combinatorial classification of toric log del Pezzo surfaces*, LMS J. Comput. Math. **13** (2010), 33-46.

[Kn] F. Knop, *Über Hilberts vierzehntes Problem für Varietäten mit Kompliziertheit eins*, Math. Z. **213** (1993), no.1, 3336.

[Ko] H. Kojima, *Logarithmic del Pezzo surfaces of rank one with unique singular points*, Japan. J. Math. (N.S.) **25** (1999), no.2, 343375.

[KM] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math. **134**, Cambridge Univ. Press, Cambridge (1998).

[LT] A. Laface and D. Testa, *Nef and semiample divisors on rational surfaces*, to appear in *Torsors, étale homotopy and applications to rational points.*

[MP] R. Miranda and U. Persson, *On extremal rational elliptic surfaces*, Math. J. **193** (1986), no.4, 537-558.

[Na] N. Nakayama, *Classification of log del Pezzo surfaces of index two*, J. Math. Sci. Univ. Tokyo **14** (2007), no.3, 293-498.

[Ni] V.V. Nikulin, *A remark on algebraic surfaces with polyhedral Mori cone*, Nagoya Math. J. **157** (2000), 73-92.

[OT] H. Ohashi and S. Taki, *K3 surfaces and log del Pezzo surfaces of index three*, Manuscripta Math. **139** (2012), no.3-4, 443471.

[Ok] S. Okawa, *On images of Mori dream spaces*, preprint, arXiv:1104.1326v1.

[OW] P. Orlik and P. Wagreich, *Algebraic surfaces with k^*-action*, Acta Math. **138** (1977), no.1-2, 43-81.

[Ot] J.C. Ottem, *On the Cox ring of $\mathbb{P}^2$ blown up in points on a line*, Math. Scand. **109** (2011), no.1, 22-30.

[Sa] F. Sakai, *Anticanonical models of rational surfaces*, Math. Ann. **269** (1984), no.3, 389-410.

[Sü] H. Süß, *Canonical divisors on T-varieties*, preprint, arXiv:0811.0626v2.

[Ta] H. Tanaka, *Minimal models and abundance for positive characteristic log surfaces*, preprint, arXiv:1201.5699v2.

[TVAV] D. Testa, A. Várilly-Alvarado, and M. Velasco, *Big rational surfaces*, Math. Ann. **351** (2011), no.1, 95-107.

[To] B. Totaro, *The cone conjecture for Calabi-Yau pairs in dimension 2*, Duke Math. J. **154** (2010), no.2, 241-263.

Department of Mathematics, Ajou University, Suwon, Korea
E-mail address: `dshwang@ajou.ac.kr`

Department of Mathematical Sciences, KAIST, Daejeon, Korea
E-mail address: `parkjh13@kaist.ac.kr`